

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property
Organization
International Bureau



(43) International Publication Date
22 April 2004 (22.04.2004)

PCT

(10) International Publication Number
WO 2004/033428 A1

(51) International Patent Classification⁷: C07D 211/58,
413/14, 409/14, 401/14, 471/04, 405/14, 417/14

(21) International Application Number:
PCT/EP2003/050697

(22) International Filing Date: 7 October 2003 (07.10.2003)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
PCT/EP02/11328 8 October 2002 (08.10.2002) EP
PCT/EP02/14836 23 December 2002 (23.12.2002) EP

(71) Applicant (for all designated States except US):
JANSSEN PHARMACEUTICA N.V. [BE/BE]; Turnhoutseweg 30, B-2340 Beerse (BE).

(72) Inventors; and

(75) Inventors/Applicants (for US only): JANSSENS, Frans, Eduard [BE/BE]; Janssen Pharmaceutica N.V., Turnhoutseweg 30, B-2340 Beerse (BE). SOMMEN, François, Maria [BE/BE]; Janssen Pharmaceutica N.V., Turnhoutseweg 30, B-2340 Beerse (BE). DE BOECK, Benoît, Christian, Albert, Ghislain [BE/BE]; Janssen Pharmaceutica N.V., Turnhoutseweg 30, B-2340 Beerse (BE). LEENAERTS, Joseph, Elisabeth [BE/BE]; Janssen Pharmaceutica N.V., Turnhoutseweg 30, B-2340 Beerse (BE). VAN ROOSBROECK, Yves, Emiel, Maria [BE/BE]; Janssen Pharmaceutica N.V., Turnhoutseweg 30, B-2340

Beerse (BE). DIELS, Gaston, Stanislas, Marcella [BE/BE]; Janssen Pharmaceutica N.V., Turnhoutseweg 30, B-2340 Beerse (BE).

(74) Common Representative: JANSSEN PHARMACEUTICA N.V.; Turnhoutseweg 30, B-2340 Beerse (BE).

(81) Designated States (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.


(84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Declarations under Rule 4.17:

- as to applicant's entitlement to apply for and be granted a patent (Rule 4.17(ii)) for all designations
- as to the applicant's entitlement to claim the priority of the earlier application (Rule 4.17(iii)) for all designations
- as to the applicant's entitlement to claim the priority of the earlier application (Rule 4.17(iii)) for all designations

[Continued on next page]

(54) Title: SUBSTITUTED 1,4-DI-PIPERIDIN-4-YL-PIPERAZINE DERIVATIVES AND THEIR USE AS NEUROKININ ANTAGONISTS



(57) Abstract: The invention concerns substituted 1,4-di-piperidin-4-yl-piperazine derivatives having neurokinin antagonistic activity, in particular NK₁ antagonistic activity, their preparation, compositions comprising them and their use as a medicine, in particular for the treatment of emesis, anxiety, depression, pain, pancreatitis and IBS. The compounds according to the invention can be represented by general Formula (I), and comprises also the pharmaceutically acceptable acid or base addition salts thereof, the stereochemically isomeric forms thereof, the N-oxide form thereof and prodrugs thereof, wherein all substituents are defined as in Claim 1. In view of their capability to antagonize the actions of neurokinins by blocking the neurokinin receptors, and in particular antagonizing the actions of substance P by blocking the NK₁ receptor, the compounds according to the invention are useful as a medicine, in particular in the prophylactic and therapeutic treatment of neurokinin-mediated conditions, such as, for instance CNS disorders, in particular depression, anxiety disorders, stress-related disorders, sleep disorders, cognitive disorders, personality disorders, schizoaffective disorders, eating disorders, neurodegenerative diseases, addiction disorders, mood disorders, sexual dysfunction, pain and other CNS-related conditions; inflammation; allergic disorders; emesis; gastrointestinal disorders, in particular irritable bowel syndrome (IBS); skin disorders; vasospastic diseases; fibrosing and collagen diseases; disorders related to immune enhancement or suppression and rheumatic diseases and body weight control.

BEST AVAILABLE COPY

WO 2004/033428 A1

82



- *of inventorship (Rule 4.17(iv)) for US only*

Published:

- *with international search report*
- *before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments*

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

-1-

SUBSTITUTED 1,4-DI-PIPERIDIN-4-YL-PIPERAZINE DERIVATIVES
AND THEIR USE AS NEUROKININ ANTAGONISTS

5 Field of the Invention

 This invention concerns 1,4-di-piperidin-4-yl-piperazine derivatives having neurokinin antagonistic activity, in particular NK₁ antagonistic activity, their preparation, compositions comprising them and their use as a medicine, in particular for the treatment of emesis, anxiety, depression, pancreatitis and IBS.

10

Background of The Invention

 Neurokinins belong to a family of short peptides that are widely distributed in the mammalian central and peripheral nervous system (Bertrand and Geppetti, *Trends Pharmacol. Sci.* 17:255-259 (1996) ; Lundberg, *Can. J. Physiol. Pharmacol.* 73:908-914 (1995) ; Maggi, *Gen. Pharmacol.* 26:911-944 (1995) ; Regoli *et al.*, *Pharmacol. Rev.* 46 (1994)). They share the common C-terminal sequence Phe-Xaa-Gly-Leu-Met-NH₂. Neurokinins released from peripheral sensory nerve endings are believed to be involved in neurogenic inflammation. In the spinal cord/central nervous system, neurokinins may play a role in pain transmission/perception and in some autonomic reflexes and behaviours. The three major neurokinins are Substance P (SP), Neurokinin A (NK_A) and Neurokinin B (NK_B) with preferential affinity for three distinct receptor subtypes, termed NK₁, NK₂, and NK₃, respectively. However, functional studies on cloned receptors suggest strong functional cross-interaction between the 3 neurokinins and their corresponding receptors (Maggi and Schwartz, *Trends Pharmacol. Sci.* 18: 351-355 (1997)).

 Species differences in structure of NK₁ receptors are responsible for species-related potency differences of NK₁ antagonists (Maggi, *Gen. Pharmacol.* 26:911-944 (1995) ; Regoli *et al.*, *Pharmacol. Rev.* 46(4):551-599 (1994)). The human NK₁ receptor closely resembles the NK₁ receptor of guinea-pigs and gerbils but differs markedly from the NK₁ receptor of rodents. The development of neurokinin antagonists has led to date to a series of peptide compounds of which might be anticipated that they are metabolically too labile to be employed as pharmaceutically active substances (Longmore J. *et al.*, *DN&P* 8(1):5-23 (1995)).

35

 The neurokinins are involved in emesis, (stress-related) anxiety states, inflammatory responses, smooth muscle contraction and pain perception. Neurokinin

-2-

antagonists are in development for indications such as emesis, anxiety and depression, irritable bowel syndrome (IBS), circadian rhythm disturbances, visceral pain, neurogenic inflammation, asthma, micturition disorders, and nociception.

5 Emesis

Nausea and vomiting are among the most distressing side effects of cancer chemotherapy. These reduce the quality of life and may cause patients to delay or refuse, potentially curative drugs (Kris *et al.*, *J. Clin. Oncol.*, 3:1379-1384 (1985)). The incidence, intensity and pattern of emesis are determined by different factors, such as the chemotherapeutic agent, dosage and route of administration. Typically, early or acute emesis starts within the first 4 h after chemotherapy administration, reaching a peak between 4 h and 10 h, and decreases by 12 to 24 h. Delayed emesis (developing after 24 h and continuing until 3-5 days post chemotherapy) is observed with most 'high-emetogenic' chemotherapeutic drugs (level 4 and 5 according to Hesketh *et al.*, *J. Clin. Oncol.* 15:103 (1997)). In humans, these 'high-emetogenic' anti-cancer treatments, including cis-platinum, induce acute emesis in > 98% and delayed emesis in 60-90% of cancer patients.

Animal models of chemotherapy such as cisplatin-induced emesis in ferrets (Rudd and Naylor, *Neuropharmacology* 33:1607-1608 (1994); Naylor and Rudd, *Cancer Surv.* 21:117-135 (1996)) have successfully predicted the clinical efficacy of the 5-HT₃ receptor antagonists. Although this discovery led to a successful therapy for the treatment of chemotherapy- and radiation-induced sickness in cancer patients, 5-HT₃ antagonists such as ondansetron and granisetron (either or not associated with dexamethasone) are effective in the control of the acute emetic phase (the first 24 h) but can only reduce the development of delayed emesis (> 24 h) with poor efficacy (De Mulder *et al.*, *Annals of Internal Medicine* 113:834-840 (1990) ; Roila, *Oncology* 50:163-167 (1993)). Despite these currently most effective treatments for the prevention of both acute and delayed emesis, still 50% of patients suffer from delayed vomiting and/or nausea (Antiemetic Subcommittee, *Annals Oncol.* 9:811-819 (1998)).

In contrast to 5-HT₃ antagonists, NK₁ antagonists such as CP-99,994 (Piedimonte *et al.*, *L. Pharmacol. Exp. Ther.* 266:270-273 (1993)) and aprepitant (also known as MK-869 or L-754,030 ; Kramer *et al.*, *Science* 281:1640-1645 (1998) ; Rupniak and Kramer, *Trends Pharmacol. Sci.* 20:1-12 (1999)) have now been shown to inhibit not only the acute but also the delayed phase of cisplatin-induced emesis in animals (Rudd *et al.*, *Br. J. Pharmacol.* 119:931-936 (1996) ; Tattersall *et al.*, *Neuropharmacology* 39:652-663 (2000)). NK₁ antagonists have also been demonstrated to reduce 'delayed'

-3-

emesis in man in the absence of concomitant therapy (Cocquyt *et al.*, *Eur. J. Cancer* 37:835-842 (2001) ; Navari *et al.*, *N. Engl. L. Med.* 340:190-195 (1999)). When administered together with dexamethasone and 5-HT₃ antagonists, moreover, NK₁ antagonists (such as MK-869 and CJ-11,974, also known as Ezlopitant) have been shown to produce additional effects in the prevention of acute emesis (Campos *et al.*, *J. Clin. Oncol.* 19:1759-1767 (2001) ; Hesketh *et al.*, *Clin. Oncol.* 17:338-343 (1999)).

Central neurokinin NK₁ receptors play a major role in the regulation of emesis. NK₁ antagonists are active against a wide variety of emetic stimuli (Watson *et al.*, *Br. J. Pharmacol.* 115:84-94 (1995) ; Tattersall *et al.*, *Neuropharmacol.* 35:1121-1129 (1996) ; Megens *et al.*, *J. Pharmacol. Exp. Ther.* 1-14 (2002)). The compounds are suggested to act by blocking central NK₁-receptors in the nucleus tractus solitarius. Apart from NK₁ antagonism, CNS penetration is thus a prerequisite for the antiemetic activity of these compounds. Loperamide-induced emesis in ferrets can be used as a fast and reliable screening model for the antiemetic activity of NK₁ antagonists. Further evaluation of their therapeutic value in the treatment of both the acute and the delayed phases of cisplatin-induced emesis has been demonstrated in the established ferret model (Rudd *et al.*, *Br. J. Pharmacol.* 119:931-936 (1994)). This model studies both 'acute' and 'delayed' emesis after cisplatin and has been validated in terms of its sensitivity to 5-HT₃ receptor antagonists, glucocorticoids (Sam *et al.*, *Eur. J. Pharmacol.* 417:231-237 (2001)) and other pharmacological challenges. It is unlikely that any future anti-emetic would find clinical acceptance unless successfully treating both the 'acute' and 'delayed' phases of emesis.

25 Depression and Anxiety

Depression is one of the most common affective disorders of modern society with a high and still increasing prevalence, particularly in the younger members of the population. The life time prevalence rates of Major depression (MDD, DSM-IV) is currently estimated to be 10-25 % for women and 5-12 % for men, whereby in about 25 % of patients the life time MDD is recurrent, without full inter-episode recovery and superimposed on dysthymic disorder. There is a high co-morbidity of depression with other mental disorders and, particularly in younger population high association with drug and alcohol abuse. In the view of the fact that depression primarily affects the population between 18-44 years of age e.g. the most productive population, it is obvious that it imposes a high burden on individuals, families and the whole society.

-4-

Among all therapeutic possibilities, the therapy with antidepressants is incontestably the most effective. A large number of antidepressants have been developed and introduced to the market in the course of the last 40 years. Nevertheless, none of the current antidepressants fulfill all criteria of an ideal drug (high therapeutic and prophylactic efficacy, rapid onset of action, completely satisfactory short- and long-term safety, simple and favorable pharmacokinetics) or is without side effects which in one or the other way limits their use in all groups and subgroups of depressed patients.

Since no treatment of the cause of depression exists at present, nor appears imminent, and no antidepressant is effective in more than 60-70 % of patients; the development of a new antidepressant which may circumvent any of the disadvantages of the available drugs is justified.

Several findings indicate involvement of SP in stress-related anxiety states. Central injection of SP induces a cardiovascular response resembling the classical "fight or flight" reaction characterized physiologically by vascular dilatation in skeletal muscles and decrease of mesenteric and renal blood flow. This cardiovascular reaction is accompanied by a behavioral response observed in rodents after noxious stimuli or stress (Culman and Unger, *Can. J. Physiol. Pharmacol.* 73:885-891 (1995)). In mice, centrally administered NK₁ agonists and antagonists are anxiogenic and anxiolytic, respectively (Teixeira *et al.*, *Eur. J. Pharmacol.* 311:7-14 (1996)). The ability of NK₁ antagonists to inhibit thumping induced by SP (or by electric shock; Ballard *et al.*, *Trends Pharmacol. Sci.* 17:255-259 (2001)) might correspond to this antidepressant / anxiolytic activity, since in gerbils thumping plays a role as an alerting or warning signal to conspecifics.

The NK₁ receptor is widely distributed throughout the limbic system and fear-processing pathways of the brain, including the amygdala, hippocampus, septum, hypothalamus, and periaqueductal grey. Additionally, substance P is released centrally in response to traumatic or noxious stimuli and substance P-associated neurotransmission may contribute to or be involved in anxiety, fear, and the emotional disturbances that accompany affective disorders such as depression and anxiety. In support of this view, changes in substance P content in discrete brain regions can be observed in response to stressful stimuli (Brodin *et al.*, *Neuropeptides* 26:253-260 (1994)).

Central injection of substance P mimetics (agonists) induces a range of defensive behavioral and cardiovascular alterations including conditioned place aversion (Elliott,

Exp. Brain. Res. 73:354-356 (1988)), potentiated acoustic startle response (Krase *et al.*, *Behav. Brain. Res.* 63:81-88 (1994)), distress vocalizations, escape behavior (Kramer *et al.*, *Science* 281:1640-1645 (1998)) and anxiety on the elevated plus maze (Aguar and Brandao, *Physiol. Behav.* 60:1183-1186 (1996)). These compounds did not modify motor performance and co-ordination on the rotarod apparatus or ambulation in an activity cage. Down-regulation of substance P biosynthesis occurs in response to the administration of known anxiolytic and antidepressant drugs (Brodin *et al.*, *Neuropeptides* 26:253-260 (1994) ; Shirayama *et al.*, *Brain. Res.* 739:70-78 (1996)). Similarly, a centrally administered NK₁ agonist-induced vocalization response in guinea-pigs can be antagonized by antidepressants such as imipramine and fluoxetine as well as L-733,060, an NK₁ antagonist. These studies provide evidence suggesting that blockade of central NK₁ receptors may inhibit psychological stress in a manner resembling antidepressants and anxiolytics (Rupniak and Kramer, *Trends Pharmacol. Sci.* 20:1-12 (1999)), but without the side effects of present medications. This is supported by data demonstrating that aprepitant is a clinically efficacious antidepressant with significant anxiolytic activity in a randomized double-blind placebo-controlled study (Kramer *et al.*, *Science* 281:1640-1645 (1998)).

Irritable bowel syndrome (IBS)

Patients with irritable bowel syndrome (IBS) experience impaired quality of life, and utilize health care resources extensively as they seek better "solutions" (including unnecessary repeated investigations or even surgery). Although these patients suffer from a 'benign' disorder (in other words, they will never die or develop significant complications), they nevertheless cause a significant economic burden by extensive health care resource utilization, and absence from work.

A reasonable number of pre-clinical publications over the role of NK₁ receptors in visceral pain has been published. Using NK₁ receptor knockout mice and NK₁ antagonists in animal models, different groups have demonstrated the important role played by the NK₁ receptor in hyperalgesia and visceral pain. The distribution of NK₁ receptors and substance P favors a major role in visceral rather than in somatic pain. Indeed more than 80% of visceral primary afferent contain substance P compared with only 25% skin afferents. NK₁ receptors are also involved in gastrointestinal motility (Tonini *et al.*, *Gastroenterol.* 120:938-945 (2001) ; Okano *et al.*, *J. Pharmacol. Exp. Ther.* 298:559-564 (2001)). Because of this dual role in both gastrointestinal motility and in nociception, NK₁ antagonists are considered to have potential to ameliorate symptoms in IBS patients.

Pancreatitis

It has been suggested that substance-P, as one of two main neuropeptides involved in neurogenic inflammation acting via the NK₁ receptor, plays an important
5 proinflammatory role in regulating the severity of acute pancreatitis and pancreatitis-associated lung injury. Impaired lung function in severe acute pancreatitis is the primary cause of morbidity and mortality in this condition. This has been clearly demonstrated using NK₁^{-/-} mice (Bhatia et al., *Proceedings of the National Academy of Sciences of the USA* 95(8):4760-4765 (1998)). Also, it has been shown that preprotachykinin-A
10 gene deletion protects mice against acute pancreatitis (Bhatia et al., *American J. of Physiology* 284(5):G380-G386(2003)). Furthermore, several studies have been published indicating that NK₁-antagonists (e.g. CP-96 345 and CP-99 994) may play a role in treating pancreatitis, e.g. in influencing the intensity, duration and frequency of pain in chronic pancreatitis patients (Maa et al., *American J. of Physiology* 279:G726-
15 G732 (2000) ; Grady et al., *British J. of Pharmacology* 130(3):505-512 (2000) ; Vera-Portocarrero et al., *Anesthesiology* 98(2):474-484 (2003) and Shrikhande et al., *Pain* 91(3):209-217 (2001)).

Background prior art


20 Compounds containing the 1-piperidin-4-yl-piperazinyl moiety were published in WO 97/16440-A1, published May 9, 1997 by Janssen Pharmaceutica N.V. for use as substance P antagonists, in WO 02/32867, published April 25, 2002 by Glaxo Group Ltd. for their special advantages as neurokinin antagonists (more specifically were disclosed 4-piperazin-1-yl-piperidine-1-carboxylic acid amide derivatives), in
25 WO 01/30348-A1, published May 03, 2001 by Janssen Pharmaceutica N.V., for use as substance P antagonists for influencing the circadian timing system, and in WO 02/062784-A1, published August 15, 2002 by Hoffmann-La Roche AG for use as neurokinin 1 antagonists.

30 The compounds of the present invention differ from the compounds of the prior art in the substitution of the piperazinyl moiety, being a substituted piperidinyl moiety as well as in their improved ability as potent, orally and centrally active neurokinin antagonists with therapeutic value, especially for the treatment of emesis, anxiety and depression, irritable bowel syndrome (IBS), circadian rhythm disturbances, visceral pain,
35 neurogenic inflammation, asthma, micturition disorders, and nociception.

Description of the Invention

-7-

The present invention relates to novel 1,4-di-piperidin-4-yl-piperazine derivatives according to the general Formula (I)



5 the pharmaceutically acceptable acid or base addition salts thereof, the stereochemically isomeric forms thereof, the *N*-oxide form thereof and prodrugs thereof, wherein :

- n is an integer, equal to 0, 1 or 2 ;
- m is an integer, equal to 1 or 2, provided that if m is 2, then n is 1 ;
- p is an integer equal to 1 or 2 ;
- 10 Q is O or NR³ ;
- X is a covalent bond or a bivalent radical of formula -O-, -S- or -NR³- ;
- each R³ independently from each other, is hydrogen or alkyl ;
- each R¹ independently from each other, is selected from the group of Ar¹, Ar¹-alkyl and di(Ar¹)-alkyl ;
- 15 q is an integer equal to 0 or 1 ;
- R² is alkyl, Ar², Ar²-alkyl, Het¹ or Het¹-alkyl ;
- Y is a covalent bond or a bivalent radical of formula -C(=O)- or -SO₂-;
- each Alk represents, independently from each other, a covalent bond; a bivalent straight or branched, saturated or unsaturated hydrocarbon radical having
- 20 from 1 to 6 carbon atoms ; or a cyclic saturated or unsaturated hydrocarbon radical having from 3 to 6 carbon atoms ; each radical optionally substituted on one or more carbon atoms with one or more alkyl, phenyl, halo, cyano, hydroxy, formyl and amino radicals ;
- L is selected from the group of hydrogen, alkyloxy, Ar³-oxy, alkyloxycarbonyl, mono- and di(alkyl)amino, mono- and di(Ar³)amino,
- 25 mono- and di(alkyloxycarbonyl)amino, Ar³, Ar³-carbonyl, Het² and Het²-carbonyl;
- Ar¹ is phenyl, optionally substituted with 1, 2 or 3 substituents each independently from each other selected from the group of halo, alkyl, cyano, aminocarbonyl and alkyloxy ;
- 30 Ar² is naphthalenyl or phenyl, each optionally substituted with 1, 2 or 3 substituents, each independently from each other, selected from the group

-8-

- of halo, nitro, amino, mono- and di(alkyl)amino, cyano, alkyl, hydroxy, alkyloxy, carboxyl, alkyloxycarbonyl, aminocarbonyl and mono- and di(alkyl)aminocarbonyl ;
- 5 Ar³ is naphthalenyl or phenyl, optionally substituted with 1, 2 or 3 substituents each independently from each other selected from the group of alkyloxy, alkyl, halo, hydroxy, pyridinyl, morpholinyl, pyrrolidinyl, imidazo[1,2-*a*]pyridinyl, morpholinylcarbonyl, pyrrolidinylcarbonyl, amino and cyano;
- 10 Het¹ is a monocyclic heterocyclic radical selected from the group of pyrrolyl, pyrazolyl, imidazolyl, furanyl, thienyl, oxazolyl, isoxazolyl, thiazolyl, thiadiazolyl, isothiazolyl, pyridinyl, pyrimidinyl, pyrazinyl and pyridazinyl; or a bicyclic heterocyclic radical selected from the group of quinolinyl, quinoxalinyl, indolyl, benzimidazolyl, benzoxazolyl, benzisoxazolyl, benzothiazolyl, benzisothiazolyl, benzofuranyl and benzothienyl ; each
- 15 heterocyclic radical may optionally be substituted on any atom by a radical selected from the group of halo and alkyl ;
- 20 Het² is a monocyclic heterocyclic radical selected from the group of pyrrolidinyl, dioxolyl, imidazolidinyl, pyrazolidinyl, piperidinyl, morpholinyl, dithianyl, thiomorpholinyl, piperazinyl, imidazolidinyl, tetrahydrofuranyl, 2H-pyrrolyl, pyrrolinyl, imidazolinyl, pyrazolinyl, pyrrolyl, imidazolyl, pyrazolyl, triazolyl, furanyl, thienyl, oxazolyl, isoxazolyl, thiazolyl, thiadiazolyl, isothiazolyl, pyridinyl, pyrimidinyl, pyrazinyl, pyridazinyl and triazinyl ; or a bicyclic heterocyclic radical selected from the group of benzopiperidinyl, quinolinyl, quinoxalinyl, indolyl, isoindolyl, chromenyl,
- 25 benzimidazolyl, imidazo[1,2-*a*]pyridinyl, benzoxazolyl, benzisoxazolyl, benzothiazolyl, benzisothiazolyl, benzofuranyl and benzothienyl ; each radical optionally substituted with one or more radicals selected from the group of Ar¹, Ar¹alkyl, halo, hydroxy, alkyl, piperidinyl, pyrrolyl, thienyl, oxo, alkyloxy, alkyloxyalkyl and alkyloxycarbonyl ; and
- 30 alkyl is a straight or branched saturated hydrocarbon radical having from 1 to 6 carbon atoms or a cyclic saturated hydrocarbon radicals having from 3 to 6 carbon atoms ; optionally substituted on one or more carbon atoms with one or more radicals selected from the group of phenyl, halo, cyano, oxo, hydroxy, formyl and amino radicals.

35

More in particular, the invention relates to a compound according to the general Formula (I), the pharmaceutically acceptable acid or base addition salts thereof, the

-9-


stereochemically isomeric forms thereof, the *N*-oxide form thereof and a prodrug thereof, wherein :

- n is 1 ;
- m is 1 ;
- 5 p is 1 ;
- Q is O ;
- X is a covalent bond ;
- each R¹ is Ar¹ or Ar¹-alkyl ;
- q is 0 or 1 ;
- 10 R² is Ar² ;
- Y is a covalent bond or a bivalent radical of formula -C(=O)- or -SO₂- ;
- each Alk represents, independently from each other, a covalent bond; a bivalent straight or branched, saturated or unsaturated hydrocarbon radical having from 1 to 6 carbon atoms ; or a cyclic saturated or unsaturated hydrocarbon radical having from 3 to 6 carbon atoms ; each radical optionally substituted on one or more carbon atoms with one or more phenyl, halo, cyano, hydroxy, formyl and amino radicals ;
- 15 L is selected from the group of hydrogen, alkyloxy, Ar³-oxy, alkyloxycarbonyl, mono- and di(alkyl)amino, mono- and di(Ar³)amino, Ar³ and Het²;
- 20 Ar¹ is phenyl, optionally substituted with 1, 2 or 3 alkyl radicals ;
- Ar² is phenyl, optionally substituted with 1, 2 or 3 alkyl radicals ;
- Ar³ is phenyl, optionally substituted with 1, 2 or 3 substituents each independently from each other selected from the group of alkyloxy, alkyl, halo, hydroxy, pyridinyl, morpholinyl, pyrrolidinyl, imidazo[1,2-*a*]pyridinyl, morpholinylcarbonyl, pyrrolidinylcarbonyl, amino and cyano ;
- 25 Het² is a monocyclic heterocyclic radical selected from the group of pyrrolidinyl, piperidinyl, morpholinyl, pyrrolyl, imidazolyl, pyrazolyl, furanyl, thienyl, isoxazolyl, thiazolyl, thiadiazolyl, pyridinyl, pyrimidinyl, pyrazinyl, and pyridazinyl ; or a bicyclic heterocyclic radical selected from the group of benzopiperidinyl, quinolinyl, quinoxalinyl, indolyl, chromenyl and benzimidazolyl ; each radical optionally substituted with one or more radicals selected from the group of Ar¹, Ar¹alkyl, halo, hydroxy, alkyl, piperidinyl, pyrrolyl, thienyl, oxo and alkyloxycarbonyl ; and
- 30 alkyl is a straight hydrocarbon radical having 1 to 6 carbon atoms, optionally substituted with one or more halo radicals ;
- 35

More in particular, the invention relates to a compound according to the general

-10-

Formula (I), the pharmaceutically acceptable acid or base addition salts thereof, the stereochemically isomeric forms thereof, the *N*-oxide form thereof and a prodrug thereof, wherein R¹ is Ar¹methyl and attached to the 2-position or R¹ is Ar¹ and attached to the 3-position, as exemplified in either of the following formulas for compounds
 5 according to Formula (I) wherein m and n are equal to 1 and Ar is an unsubstituted phenyl. Preferably, Ar¹methyl is an unsubstituted benzyl radical.



More in particular, the invention relates to a compound according to the general
 10 Formula (I), the pharmaceutically acceptable acid or base addition salts thereof, the stereochemically isomeric forms thereof, the *N*-oxide form thereof and a prodrug thereof, wherein the R²-X-C(=Q)- moiety is 3,5-di-(trifluoromethyl) phenylcarbonyl.

In the framework of this application, alkyl is defined as a monovalent straight or
 15 branched saturated hydrocarbon radical having from 1 to 6 carbon atoms, for example methyl, ethyl, propyl, butyl, 1-methylpropyl, 1,1-dimethylethyl, pentyl, hexyl ; alkyl further defines a monovalent cyclic saturated hydrocarbon radical having from 3 to 6 carbon atoms, for example cyclopropyl, methylcyclopropyl, cyclobutyl, cyclopentyl and cyclohexyl. The definition of alkyl also comprises an alkyl radical that is optionally
 20 substituted on one or more carbon atoms with one or more phenyl, halo, cyano, oxo, hydroxy, formyl and amino radicals, for example hydroxyalkyl, in particular hydroxymethyl and hydroxyethyl and polyhaloalkyl, in particular difluoromethyl and trifluoromethyl.

25 In the framework of this application, halo is generic to fluoro, chloro, bromo and iodo.

In the framework of this application, with "compounds according to the invention" is meant a compound according to the general Formula (I), the pharmaceutically
 30 acceptable acid or base addition salts thereof, the stereochemically isomeric forms thereof, the *N*-oxide form thereof and a prodrug thereof.

In the framework of this application, especially in the moiety $\text{Alk}^a\text{-Y-Alk}^b$ in Formula (I), when two or more consecutive elements of said moiety denote a covalent bond, then a single covalent bond is denoted. For example, when Alk^a and Y denote both a covalent bond and Alk^b is $-\text{CH}_2-$, then the moiety $\text{Alk}^a\text{-Y-Alk}^b$ denotes $-\text{CH}_2-$.

The pharmaceutically acceptable salts are defined to comprise the therapeutically active non-toxic acid addition salts forms that the compounds according to Formula (I) are able to form. Said salts can be obtained by treating the base form of the compounds according to Formula (I) with appropriate acids, for example inorganic acids, for example hydrohalic acid, in particular hydrochloric acid, hydrobromic acid, sulfuric acid, nitric acid and phosphoric acid; organic acids, for example acetic acid, hydroxyacetic acid, propanoic acid, lactic acid, pyruvic acid, oxalic acid, malonic acid, succinic acid, maleic acid, fumaric acid, malic acid, tartaric acid, citric acid, methanesulfonic acid, ethanesulfonic acid, benzenesulfonic acid, p-toluenesulfonic acid, cyclamic acid, salicylic acid, p-aminosalicylic acid and pamoic acid. Particularly preferred is fumaric acid.

The compounds according to Formula (I) containing acidic protons may also be converted into their therapeutically active non-toxic metal or amine addition salts forms by treatment with appropriate organic and inorganic bases. Appropriate base salts forms comprise, for example, the ammonium salts, the alkaline and earth alkaline metal salts, in particular lithium, sodium, potassium, magnesium and calcium salts, salts with organic bases, e.g. the benzathine, *N*-methyl-D-glucamine, hybamine salts, and salts with amino acids, for example arginine and lysine.

Conversely, said salts forms can be converted into the free forms by treatment with an appropriate base or acid.

The term addition salt as used in the framework of this application also comprises the solvates that the compounds according to Formula (I) as well as the salts thereof, are able to form. Such solvates are, for example, hydrates and alcoholates.

The *N*-oxide forms of the compounds according to Formula (I) are meant to comprise those compounds of Formula (I) wherein one or several nitrogen atoms are oxidized to the so-called *N*-oxide, particularly those *N*-oxides wherein one or more tertiary nitrogens (e.g. of the piperazinyl or piperidinyl radical) are *N*-oxidized. Such

-12-

N-oxides can easily be obtained by a skilled person without any inventive skills and they are obvious alternatives for the compounds according to Formula (I) since these compounds are metabolites, which are formed by oxidation in the human body upon uptake. As is generally known, oxidation is normally the first step involved in drug metabolism (Textbook of Organic Medicinal and Pharmaceutical Chemistry, 1977, pages 70- 75). As is also generally known, the metabolite form of a compound can also be administered to a human instead of the compound per se, with much the same effects.

The compounds according to the invention possess at least 2 oxydizable nitrogens (tertiary amines moieties). It is therefore highly likely that *N*-oxides are to form in the human metabolism.

The compounds of Formula (I) may be converted to the corresponding *N*-oxide forms following art-known procedures for converting a trivalent nitrogen into its *N*-oxide form. Said *N*-oxidation reaction may generally be carried out by reacting the starting material of Formula (I) with an appropriate organic or inorganic peroxide. Appropriate inorganic peroxides comprise, for example, hydrogen peroxide, alkali metal or earth alkaline metal peroxides, e.g. sodium peroxide, potassium peroxide; appropriate organic peroxides may comprise peroxy acids such as, for example, benzenecarboperoxoic acid or halo substituted benzenecarboperoxoic acid, e.g. 3-chlorobenzenecarboperoxoic acid, peroxyalkanoic acids, e.g. peroxyacetic acid, alkylhydroperoxides, e.g. *tert*-butyl hydroperoxide. Suitable solvents are, for example, water, lower alkanols, e.g. ethanol and the like, hydrocarbons, e.g. toluene, ketones, e.g. 2-butanone, halogenated hydrocarbons, e.g. dichloromethane, and mixtures of such solvents.

The term "stereochemically isomeric forms" as used hereinbefore defines all the possible isomeric forms that the compounds of Formula (I) may possess. Unless otherwise mentioned or indicated, the chemical designation of compounds denotes the mixture of all possible stereochemically isomeric forms, said mixtures containing all diastereomers and enantiomers of the basic molecular structure. More in particular, stereogenic centers may have the R- or S-configuration; substituents on bivalent cyclic (partially) saturated radicals may have either the cis- or trans-configuration. Compounds encompassing double bonds can have an E or Z-stereochemistry at said double bond. Stereochemically isomeric forms of the compounds of Formula (I) are obviously intended to be embraced within the scope of this invention.

Following CAS nomenclature conventions, when two stereogenic centers of known absolute configuration are present in a molecule, an *R* or *S* descriptor is assigned (based on Cahn-Ingold-Prelog sequence rule) to the lowest-numbered chiral center, the reference center. The configuration of the second stereogenic center is indicated using relative descriptors [*R**,*R**] or [*R**,*S**], where *R** is always specified as the reference center and [*R**,*R**] indicates centers with the same chirality and [*R**,*S**] indicates centers of unlike chirality. For example, if the lowest-numbered chiral center in the molecule has an *S* configuration and the second center is *R*, the stereo descriptor would be specified as *S*-[*R**,*S**]. If "α" and "β" are used : the position of the highest priority substituent on the asymmetric carbon atom in the ring system having the lowest ring number, is arbitrarily always in the "α" position of the mean plane determined by the ring system. The position of the highest priority substituent on the other asymmetric carbon atom in the ring system (hydrogen atom in compounds according to Formula (I)) relative to the position of the highest priority substituent on the reference atom is denominated "α", if it is on the same side of the mean plane determined by the ring system, or "β", if it is on the other side of the mean plane determined by the ring system.


Compounds according to Formula (I) and some of the intermediate compounds have at least two stereogenic centers in their structure, namely at the 2-position of the piperidinyl-moiety (*R* or *S*) and at the 4-position, where the attached radical may be either in the *cis* or *trans* position with respect to the radical at the 2-position on the piperidinyl-moiety.

The invention also comprises derivative compounds (usually called "pro-drugs") of the pharmacologically-active compounds according to the invention, which are degraded *in vivo* to yield the compounds according to the invention. Pro-drugs are usually (but not always) of lower potency at the target receptor than the compounds to which they are degraded. Pro-drugs are particularly useful when the desired compound has chemical or physical properties that make its administration difficult or inefficient. For example, the desired compound may be only poorly soluble, it may be poorly transported across the mucosal epithelium, or it may have an undesirably short plasma half-life. Further discussion on pro-drugs may be found in Stella, V. J. *et al.*; "Prodrugs", *Drug Delivery Systems*, 1985, pp. 112-176, and *Drugs*, 1985, 29, pp. 455-473.

Pro-drugs forms of the pharmacologically-active compounds according to the invention will generally be compounds according to Formula (I), the pharmaceutically

-14-

acceptable acid or base addition salts thereof, the stereochemically isomeric forms thereof and the *N*-oxide form thereof, having an acid group which is esterified or amidated. Included in such esterified acid groups are groups of the formula $-\text{COOR}^x$, where R^x is a C_{1-6} alkyl, phenyl, benzyl or one of the following groups :



Amidated groups include groups of the formula $-\text{CONR}^y\text{R}^z$, wherein R^y is H, C_{1-6} alkyl, phenyl or benzyl and R^z is $-\text{OH}$, H, C_{1-6} alkyl, phenyl or benzyl. Compounds according to the invention having an amino group may be derivatised with a ketone or an aldehyde such as formaldehyde to form a Mannich base. This base will hydrolyze with first order kinetics in aqueous solution.

The compounds of Formula (I) as prepared in the processes described below may be synthesized in the form of racemic mixtures of enantiomers that can be separated from one another following art-known resolution procedures. The racemic compounds of Formula (I) may be converted into the corresponding diastereomeric salt forms by reaction with a suitable chiral acid. Said diastereomeric salt forms are subsequently separated, for example, by selective or fractional crystallization and the enantiomers are liberated therefrom by alkali. An alternative manner of separating the enantiomeric forms of the compounds of Formula (I) involves liquid chromatography using a chiral stationary phase. Said pure stereochemically isomeric forms may also be derived from the corresponding pure stereochemically isomeric forms of the appropriate starting materials, provided that the reaction occurs stereospecifically. Preferably if a specific stereoisomer is desired, said compound would be synthesized by stereospecific methods of preparation. These methods will advantageously employ enantiomerically pure starting materials.

Pharmacology

Substance P and other neurokinins are involved in a variety of biological actions such as pain transmission (nociception), neurogenic inflammation, smooth muscle contraction, plasma protein extravasation, vasodilation, secretion, mast cell
5 degranulation, and also in activation of the immune system. A number of diseases are deemed to be engendered by activation of neurokinin receptors, in particular the NK₁ receptor, by excessive release of substance P and other neurokinins in particular cells such as cells in the neuronal plexi of the gastrointestinal tract, unmyelinated primary sensory afferent neurons, sympathetic and parasympathetic neurons and nonneuronal
10 cell types (DN&P 8(1):5-23 (1995) and Longmore J. *et al.*, "Neurokinin Receptors" *Pharmacological Reviews* 46(4):551-599 (1994)).

The compounds of the present invention are potent inhibitors of neurokinin-mediated effects, in particular those mediated via the NK₁ receptor, and may therefore
15 be described as neurokinin antagonists or tachykinin antagonists, especially as substance P antagonists, as indicated *in vitro* by the antagonism of substance P-induced relaxation of pig coronary arteries which is described hereinafter. The binding affinity of the present compounds for the human, guinea-pig and gerbil neurokinin receptors may be determined *in vitro* in a receptor binding test using ³H-substance-P as radioligand. The
20 subject compounds also show substance-P antagonistic activity *in vivo* as may be evidenced by, for instance, the antagonism of substance P-induced plasma extravasation in guinea-pigs, or the antagonism of drug-induced emesis in ferrets (Watson *et al.*, *Br. J. Pharmacol.* 115:84-94 (1995)).

25 In view of their capability to antagonize the actions of neurokinins by blocking the neurokinin receptors, and in particular antagonizing the actions of substance P by blocking the NK₁ receptor, the compounds according to the invention are useful as a medicine, in particular in the prophylactic and therapeutic treatment of neurokinin-mediated conditions.

30 The invention therefore relates to a compound according to the general Formula (I), the pharmaceutically acceptable acid or base addition salts thereof, the stereochemically isomeric forms thereof, the *N*-oxide form thereof and prodrugs thereof, for use as a medicine.

35

-16-

The invention also relates to the use of a compound according to any one of claims 1-3 for the manufacture of a medicament for treating, either prophylactic or therapeutic or both, neurokinin mediated conditions.

5 The compounds according to the invention are useful in the treatment of CNS disorders, in particular depression, anxiety disorders, stress-related disorders, sleep disorders, cognitive disorders, personality disorders, schizoaffective disorders, eating disorders, neurodegenerative diseases, addiction disorders, mood disorders, sexual dysfunction, pain and other CNS-related conditions ; inflammation ; allergic disorders ;
10 emesis ; gastrointestinal disorders, in particular irritable bowel syndrome (IBS); skin disorders ; vasospastic diseases ; fibrosing and collagen diseases ; disorders related to immune enhancement or suppression and rheumatic diseases and body weight control.

 In particular, the compounds according to the invention are useful in the
15 treatment or prevention of depression including but not limited to major depressive disorders including bipolar depression ; unipolar depression ; single or recurrent major depressive episodes with or without psychotic features, catatonic features, melancholic features, atypical features or postpartum onset, and, in the case of recurrent episodes, with or without seasonal pattern. Other mood disorders encompassed within the term
20 "major depressive disorder" include dysthymic disorder with early or late onset and with or without atypical features, bipolar I disorder, bipolar II disorder, cyclothymic disorder, recurrent brief depressive disorder, mixed affective disorder, neurotic depression, post traumatic stress disorder and social phobia ; dementia of the Alzheimer's type with early or late onset, with depressed mood ; vascular dementia with
25 depressed mood ; substance-induced mood disorders such as mood disorders induced by alcohol, amphetamines, cocaine, hallucinogens, inhalants, opioids, phencyclidine, sedatives, hypnotics, anxiolytics and other substances ; schizoaffective disorder of the depressed type ; and adjustment disorder with depressed mood. Major depressive disorders may also result from a general medical condition including, but not limited to,
30 myocardial infarction, diabetes, miscarriage or abortion, etc.

 In particular, the compounds according to the invention are useful in the treatment or prevention of anxiety disorders, including but not limited to panic attack ; agoraphobia ; panic disorder without agoraphobia ; agoraphobia without history of
35 panic disorder ; specific phobia ; social phobia ; obsessive-compulsive disorder ; post-traumatic stress disorder ; acute stress disorder ; generalized anxiety disorder ; anxiety

-17-

disorder due to a general medical condition ; substance-induced anxiety disorder ; and anxiety disorder not otherwise specified.

In particular, the compounds according to the invention are useful in the treatment or prevention of stress-related disorders associated with depression and/or anxiety, including but not limited to acute stress reaction ; adjustment disorders, such as brief depressive reaction, prolonged depressive reaction, mixed anxiety and depressive reaction, adjustment disorder with predominant disturbance of other emotions, adjustment disorder with predominant disturbance of conduct, adjustment disorder with mixed disturbance of emotions and conduct and adjustment disorders with other specified predominant symptoms ; and other reactions to severe stress.

In particular, the compounds according to the invention are useful in the treatment or prevention of sleep disorders, including but not limited to dysomnia and/or parasomnias as primary sleep disorders ; insomnia ; sleep apnea ; narcolepsy ; circadian rhythms disorders ; sleep disorders related to another mental disorder ; sleep disorder due to a general medical condition ; and substance-induced sleep disorder.

In particular, the compounds according to the invention are useful in the treatment or prevention of cognitive disorders, including but not limited to dementia ; amnesic disorders and cognitive disorders not otherwise specified, especially dementia caused by degenerative disorders, lesions, trauma, infections, vascular disorders, toxins, anoxia, vitamin deficiency or endocrinic disorders ; dementia of the Alzheimer's type, with early or late onset, with depressed mood ; AIDS-associated dementia or amnesic disorders caused by alcohol or other causes of thiamin deficiency, bilateral temporal lobe damage due to Herpes simplex encephalitis and other limbic encephalitis, neuronal loss secondary to anoxia / hypoglycemia / severe convulsions and surgery, degenerative disorders, vascular disorders or pathology around ventricle III. Furthermore, the compounds according to the invention are also useful as memory and/or cognition enhancers in healthy humans with no cognitive and/or memory deficit.

In particular, the compounds according to the invention are useful in the treatment or prevention of personality disorders, including but not limited to paranoid personality disorder ; schizoid personality disorder ; schizotypal personality disorder ; antisocial personality disorder ; borderline personality disorder ; histrionic personality disorder ; narcissistic personality disorder ; avoidant personality disorder ; dependent personality

-18-

disorder ; obsessive-compulsive personality disorder and personality disorder not otherwise specified.

In particular, the compounds according to the invention are useful in the treatment
5 or prevention of schizoaffective disorders resulting from various causes, including
schizoaffective disorders of the manic type, of the depressive type, of mixed type ;
paranoid, disorganized, catatonic, undifferentiated and residual schizophrenia ;
schizophreniform disorder ; schizoaffective disorder ; delusional disorder ; brief
psychotic disorder ; shared psychotic disorder ; substance-induced psychotic disorder ;
10 and psychotic disorder not otherwise specified.

In particular, the compounds according to the invention are also useful in the
treatment or prevention of eating disorders, including anorexia nervosa ; atypical
anorexia nervosa ; bulimia nervosa ; atypical bulimia nervosa ; overeating associated
15 with other psychological disturbances ; vomiting associated with other psychological
disturbances ; and non-specified eating disorders.

In particular, the compounds according to the invention are also useful in the
treatment or prevention of neurodegenerative diseases, including but not limited to
20 Alzheimer's disease ; Huntington's chorea ; Creutzfeld-Jacob disease ; Pick's disease ;
demyelinating disorders, such as multiple sclerosis and ALS ; other neuropathies and
neuralgia ; multiple sclerosis ; amyotropical lateral sclerosis ; stroke and head trauma.

In particular, the compounds according to the invention are also useful in the
25 treatment or prevention of addiction disorders, including but not limited to substance
dependence or abuse with or without physiological dependence, particularly where the
substance is alcohol, amphetamines, amphetamine-like substances, caffeine, cocaine,
hallucinogens, inhalants, nicotine, opioids (such as cannabis, heroin and morphine),
phencyclidine, phencyclidine-like compounds, sedative-hypnotics, benzodiazepines
30 and/or other substances, particularly useful for treating withdrawal from the above
substances and alcohol withdrawal delirium.

In particular, the compounds according to the invention are also useful in the
treatment or prevention of mood disorders induced particularly by alcohol,
35 amphetamines, caffeine, cannabis, cocaine, hallucinogens, inhalants, nicotine, opioids,
phencyclidine, sedatives, hypnotics, anxiolytics and other substances.

-19-

In particular, the compounds according to the invention are also useful in the treatment or prevention of sexual dysfunction, including but not limited to sexual desire disorders ; sexual arousal disorders ; orgasmic disorders ; sexual pain disorders ; sexual dysfunction due to a general medical condition ; substance-induced sexual dysfunction
5 and sexual dysfunction not otherwise specified.

In particular, the compounds according to the invention are also useful in the treatment or prevention of pain, including but not limited to traumatic pain such as postoperative pain ; traumatic avulsion pain such as brachial plexus ; chronic pain such
10 pancreatitis induced chronic pain or arthritic pain such as occurring in osteo- rheumatoid or psoriatic arthritis ; neuropathic pain such as post-herpetic neuralgia, trigeminal neuralgia, segmental or intercostal neuralgia, fibromyalgia, causalgia, peripheral neuropathy, diabetic neuropathy, chemotherapy-induced neuropathy, AIDS related neuropathy, occipital neuralgia, geniculate neuralgia, glossopharyngeal neuralgia, reflex
15 sympathetic dystrophy and phantom limb pain ; various forms of headache such as migraine, acute or chronic tension headache, temporomandibular pain, maxillary sinus pain and cluster headache ; odontalgia ; cancer pain ; visceral pain ; gastrointestinal pain ; nerve entrapment pain ; sport's injury pain ; dysmennorrhoea ; menstrual pain ; meningitis ; arachnoiditis ; musculoskeletal pain ; low back pain such as spinal stenosis,
20 prolapsed disc, sciatica, angina, ankylosing spondylitis ; gout ; burns ; scar pain ; itch ; and thalamic pain such as post stroke thalamic pain.

In particular, the compounds according to the invention are also useful in the treatment or prevention of the following other CNS-related conditions : akinesia,
25 akinetic-rigid syndromes, dyskinesia and medication-induced parkinsonism, Gilles de la Tourette syndrome and its symptoms, tremor, chorea, myoclonus, tics and dystonia, attention-deficit / hyperactivity disorder (ADHD), Parkinson's disease, drug-induced Parkinsonism, post-encephalitic Parkinsonism, progressive supranuclear palsy, multiple system atrophy, corticobasal degeneration, parkinsonism-ALS dementia complex and
30 basal ganglia calcification, behavioral disturbances and conduct disorders in dementia and the mentally retarded, including restlessness and agitation, extra-pyramidal movement disorders , Down's syndrome and Akathisia.

In particular, the compounds according to the invention are also useful in the
35 treatment or prevention of inflammation, including but not limited to inflammatory conditions in asthma, influenza, chronic bronchitis and rheumatoid arthritis ; inflammatory conditions in the gastrointestinal tract such as, but not limited to Crohn's

-20-

disease, ulcerative colitis, inflammatory bowel disease and non-steroidal anti-inflammatory drug induced damage ; inflammatory conditions of the skin such as herpes and eczema ; inflammatory conditions of the bladder such as cystitis and urge incontinence ; and eye and dental inflammation and pancreatitis, in particular chronic and acute pancreatitis.

In particular, the compounds according to the invention are also useful in the treatment or prevention of allergic disorders, including but not limited to allergic disorders of the skin such as but not limited to urticaria ; and allergic disorders of the airways such as but not limited to rhinitis.

In particular, the compounds according to the invention are also useful in the treatment or prevention of emesis, i.e. nausea, retching and vomiting, including but not limited to acute emesis, delayed emesis and anticipatory emesis ; emesis induced by drugs such as cancer chemotherapeutic agents such as alkylating agents, for example cyclophosphamide, carmustine, lomustine and chlorambucil ; cytotoxic antibiotics, for example dactinomycin, doxorubicin, mitomycin-C and bleomycin ; anti-metabolites, for example cytarabine, methotrexate and 5-fluorouracil ; vinca alkaloids, for example etoposide, vinblastine and vincristine ; and other drugs such as cisplatin, dacarbazine, procarbazine and hydroxyurea ; and combinations thereof ; radiation sickness ; radiation therapy, such as in the treatment of cancer ; poisons ; toxins such as toxins caused by metabolic disorders or by infection, such as gastritis, or released during bacterial or viral gastrointestinal infection ; pregnancy ; vestibular disorders, such as motion sickness, vertigo, dizziness and Meniere's disease ; post-operative sickness ; gastrointestinal obstruction ; reduced gastrointestinal motility ; visceral pain, such as myocardial infarction or peritonitis ; migraine ; increased intracranial pressure ; decreased intracranial pressure (such as altitude sickness) ; opioid analgesics, such as morphine ; gastro-oesophageal reflux disease ; acid indigestion ; over-indulgence of food or drink ; acid stomach ; sour stomach ; waterbrash/regurgitation ; heartburn, such as episodic heartburn, nocturnal heartburn and meal induced heartburn ; and dyspepsia.

In particular, the compounds according to the invention are also useful in the treatment or prevention of gastrointestinal disorders, including but not limited to irritable bowel syndrome (IBS), skin disorders such as psoriasis, pruritis and sunburn ; vasospastic diseases such as angina, vascular headache and Reynaud's disease, cerebral ischaemia such as cerebral vasospasm following subarachnoid haemorrhage ; fibrosing and collagen diseases such as scleroderma and eosinophilic fascioliasis ; disorders

related to immune enhancement or suppression such as systemic lupus erythematosus and rheumatic diseases such as fibrositis ; cough ; and body weight control, including obesity.

5 The present invention also relates to a method for the treatment and/or prophylaxis of neurokinin-mediated diseases, in particular for the treatment and/or prophylaxis of depression, anxiety disorders, emesis and irritable bowel syndrome (IBS) comprising administering to a human in need of such administration an effective amount of a compound according to the invention, in particular according to Formula (I), the
10 pharmaceutically acceptable acid or base addition salts thereof, the stereochemically isomeric forms thereof, the *N*-oxide form thereof, as well as the pro-drugs thereof.

 The invention also relates to a pharmaceutical composition comprising a pharmaceutically acceptable carrier and, as active ingredient, a therapeutically effective
15 amount of a compound according to the invention, in particular a compound according to Formula (I), the pharmaceutically acceptable acid or base addition salts thereof, the stereochemically isomeric forms thereof, the *N*-oxide form thereof and a prodrug thereof

 The compounds according to the invention, in particular the compounds according to Formula (I), the pharmaceutically acceptable acid or base addition salts thereof, the stereochemically isomeric forms thereof, the *N*-oxide form thereof and the prodrugs thereof, or any subgroup or combination thereof may be formulated into various pharmaceutical forms for administration purposes. As appropriate compositions there may be cited all compositions usually employed for systemically administering drugs.
20 To prepare the pharmaceutical compositions of this invention, an effective amount of the particular compound, optionally in addition salt form, as the active ingredient is combined in intimate admixture with a pharmaceutically acceptable carrier, which carrier may take a wide variety of forms depending on the form of preparation desired for administration. These pharmaceutical compositions are desirable in unitary dosage form suitable, in particular, for administration orally, rectally, percutaneously, by parenteral
25 injection or by inhalation. For example, in preparing the compositions in oral dosage form, any of the usual pharmaceutical media may be employed such as, for example, water, glycols, oils, alcohols and the like in the case of oral liquid preparations such as suspensions, syrups, elixirs, emulsions and solutions; or solid carriers such as starches,
30 sugars, kaolin, diluents, lubricants, binders, disintegrating agents and the like in the case of powders, pills, capsules and tablets. Because of their ease in administration, tablets and capsules represent the most advantageous oral dosage unit forms in which case solid

pharmaceutical carriers are obviously employed. For parenteral compositions, the carrier will usually comprise sterile water, at least in large part, though other ingredients, for example, to aid solubility, may be included. Injectable solutions, for example, may be prepared in which the carrier comprises saline solution, glucose solution or a mixture of saline and glucose solution. Injectable suspensions may also be prepared in which case appropriate liquid carriers, suspending agents and the like may be employed. Also included are solid form preparations that are intended to be converted, shortly before use, to liquid form preparations. In the compositions suitable for percutaneous administration, the carrier optionally comprises a penetration enhancing agent and/or a suitable wetting agent, optionally combined with suitable additives of any nature in minor proportions, which additives do not introduce a significant deleterious effect on the skin. Said additives may facilitate the administration to the skin and/or may be helpful for preparing the desired compositions. These compositions may be administered in various ways, e.g., as a transdermal patch, as a spot-on, as an ointment.

It is especially advantageous to formulate the aforementioned pharmaceutical compositions in unit dosage form for ease of administration and uniformity of dosage. Unit dosage form as used herein refers to physically discrete units suitable as unitary dosages, each unit containing a predetermined quantity of active ingredient calculated to produce the desired therapeutic effect in association with the required pharmaceutical carrier. Examples of such unit dosage forms are tablets (including scored or coated tablets), capsules, pills, powder packets, wafers, suppositories, injectable solutions or suspensions and the like, and segregated multiples thereof.

Since the compounds according to the invention are potent orally administrable NK₁ antagonists, pharmaceutical compositions comprising said compounds for administration orally are especially advantageous.


Preparation

The compounds according to the invention can generally be prepared by a succession of steps, each of which is known to the skilled person.

The compounds of Formula (I) are conveniently prepared by reductively *N*-alkylating an intermediate of Formula (II) wherein R¹, R², X, Q, m, n and p are defined as in Formula (I), with a *N*-substituted piperidinon of Formula (III) wherein R¹, Alk, Y, L and q are defined as in Formula (I). Said reductive *N*-alkylation may be performed in a reaction-inert solvent such as, for example, dichloromethane, ethanol or toluene or a

-23-

mixture thereof, and in the presence of an appropriate reducing agent such as, for example, a borohydride, e.g. sodium borohydride, sodium cyanoborohydride or triacetoxy borohydride. In case a borohydride is used as a reducing agent, it may be convenient to use a complex-forming agent such as, for example, titanium(IV)-isopropylate as described in J. Org. Chem, 1990, 55, 2552-2554. Using said complex-forming agent may also result in an improved *cis/trans* ratio in favor of the *trans* isomer. It may also be convenient to use hydrogen as a reducing agent in combination with a suitable catalyst such as, for example, palladium-on-charcoal or platinum-on-charcoal. In case hydrogen is used as reducing agent, it may be advantageous to add a dehydrating agent to the reaction mixture such as, for example, aluminium *tert*-butoxide. In order to prevent the undesired further hydrogenation of certain functional groups in the reactants and the reaction products, it may also be advantageous to add an appropriate catalyst-poison to the reaction mixture, e.g., thiophene or quinoline-sulphur. Stirring and optionally elevated temperatures and/or pressure may enhance the rate of the reaction.




In this and the following preparations, the reaction products may be isolated from the reaction medium and, if necessary, further purified according to methodologies generally known in the art such as, for example, extraction, crystallization, trituration and chromatography.


Especially advantage is the preparation of a compound according to the invention according to the previous reaction scheme in which the Alk-Y-Alk-L-moiety is benzyl, thus giving rise to a compound according to Formula (I) in which the Alk-Y-Alk-L-moiety is benzyl. Said compound is pharmacological active and can be converted into a compound according to the invention in which the Alk-Y-Alk-L-moiety is hydrogen by reductive hydrogenation using e.g. hydrogen as a reducing agent in combination with a suitable catalyst such as, for example, palladium-on-charcoal or platinum-on-charcoal. The resulting compound according to the invention can then be converted into other compounds according to the invention by art-known transformations, e.g. acylation and alkylation.

-24-

In particular, the compounds of Formula (I^a) can be prepared by reacting a final compound of Formula (I') wherein R¹, R², X, Q, m, n, p and q are defined as in Formula (I) with an acyl compound of Formula (V) wherein Alk and L are defined as in Formula (I) and W¹ is an appropriate leaving group such as, for example, a halo, e.g. chloro or bromo, or a sulfonyloxy leaving group, e.g. methanesulfonyloxy or benzenesulfonyloxy. The reaction can be performed in a reaction-inert solvent such as, for example, a chlorinated hydrocarbon, e.g. dichloromethane, an alcohol, e.g. ethanol, or a ketone, e.g. methyl isobutylketone, and in the presence of a suitable base such as, for example, sodium carbonate, sodium hydrogen carbonate or triethylamine. Stirring may enhance the rate of the reaction. The reaction may conveniently be carried out at a temperature ranging between room temperature and reflux temperature.




Alternatively, the compounds of Formula (I^a) can also be prepared by reacting a final compound of Formula (I') wherein R¹, R², X, Q, m, n, p and q are defined as in Formula (I) with a carboxylic acid of Formula (VI) wherein Alk and L are defined as in Formula (I) (base-catalyzed nucleophilic addition reaction). The reaction can be performed in a reaction-inert solvent such as, for example, a chlorinated hydrocarbon, e.g. dichloromethane, an alcohol, e.g. ethanol, or a ketone, e.g. methyl isobutylketone, and in the presence of a suitable base such as, for example, sodium carbonate, sodium hydrogen carbonate or triethylamine. Stirring may enhance the rate of the reaction. The reaction may conveniently be carried at a temperature ranging between room temperature and reflux temperature.




The above reaction may also be carried out under equivalent conditions with the carboxylic ester of the carboxylic acid of Formula (VI).

-25-

In particular, the compounds of Formula (I^b) can be prepared by reacting a final compound of Formula (I') wherein R¹, R², X, Q, m, n, p and q are defined as in Formula (I) with a keto-compound of Formula (VII) wherein W² is an appropriate leaving group such as, for example, a halogen, e.g. chloro or bromo, or a sulfonyloxy leaving group, e.g. methanesulfonyloxy or benzenesulfonyloxy. The reaction can be performed in a reaction-inert solvent such as, for example, a chlorinated hydrocarbon, e.g. dichloromethane, an alcohol, e.g. ethanol, or a ketone, e.g. methyl isobutylketone, and in the presence of a suitable base such as, for example, sodium carbonate, sodium hydrogen carbonate or triethylamine. Stirring may enhance the rate of the reaction. The reaction may conveniently be carried at a temperature ranging between room temperature and reflux temperature.




The compounds of Formula (I^c) can be prepared by reductive amination/alkylation of a final compound of Formula (I') wherein R¹, R², X, Q, m, n, p and q are defined as in Formula (I) with a compound of Formula (VIII) wherein Alk and L are defined as in Formula (I) and W³ is an appropriate leaving group such as, for example, a halogen, e.g. chloro or bromo, or a sulfonyloxy leaving group, e.g. methanesulfonyloxy or benzenesulfonyloxy. The reaction can be performed in a reaction-inert solvent such as, for example, a chlorinated hydrocarbon, e.g. dichloromethane, an alcohol, e.g. ethanol, or a ketone, e.g. methyl isobutylketone, and in the presence of a suitable base such as, for example, sodium carbonate, sodium hydrogen carbonate or triethylamine. Stirring may enhance the rate of the reaction. The reaction may conveniently be carried at a temperature ranging between room temperature and reflux temperature.



-26-

The starting materials and some of the intermediates are known compounds and are commercially available or may be prepared according to conventional reaction procedures generally known in the art. For example, intermediates of formula (II) may be prepared by reductively *N*-alkylating an intermediate of formula (IX) with an intermediate of formula (X) in which W^4 is a benzyl radical, after which the compound according to Formula (X) is subsequently reduced to yield an intermediate compound according to Formula (II). Said reductive *N*-alkylation may be performed in a reaction-inert solvent such as, for example, dichloromethane, ethanol, toluene or a mixture thereof, and in the presence of an appropriate reducing agent such as, for example, a borohydride, e.g. sodium borohydride, sodium cyanoborohydride or triacetoxy borohydride. In case a borohydride is used as a reducing agent, it may be convenient to use a complex-forming agent such as, for example, titanium(IV)isopropylate as described in J. Org. Chem, 1990, 55, 2552-2554. Using said complex-forming agent may also result in an improved *cis/trans* ratio in favor of the *trans* isomer. It may also be convenient to use hydrogen as a reducing agent in combination with a suitable catalyst such as, for example, palladium-on-charcoal or platinum-on-charcoal. In case hydrogen is used as reducing agent, it may be advantageous to add a dehydrating agent to the reaction mixture such as, for example, aluminium *tert*-butoxide. In order to prevent the undesired further hydrogenation of certain functional groups in the reactants and the reaction products, it may also be advantageous to add an appropriate catalyst-poison to the reaction mixture, e.g., thiophene or quinoline-sulphur. Stirring and optionally elevated temperatures and/or pressure may enhance the rate of the reaction.



The preparation of these and other intermediates is described in WO 97/16440-A1, published May 9, 1997 by Janssen Pharmaceutica N.V, which is disclosed herein by reference as well as in other publications mentioned in WO 97/16440-A1, such as, e.g. EP-0,532,456-A.

Compounds according to the invention may be converted into each other following art-known transformation reactions, such as illustrated further.

-27-

More specifically, compounds of Formula (XIII), wherein A is an aryl or heteroaryl, Z may be any moiety, preferably a moiety Z¹ as defined below, Het is an unsaturated heteroaryl and r is an integer ranging from 1 to a number equal to the number of available carbon atoms in the aryl or heteroaryl-moiety A, e.g. 5 in phenyl and 4 in pyrrolyl, may be obtained by a novel type of Heck-reaction wherein a compound of Formula (XI), wherein Z, A and r are as defined in Formula (XIII) and Hal is a halogen, thus comprising an active or non-active halo-substituted aryl or halo-substituted heteroaryl, more preferably a mono or polysubstituted bromo- and/or iodoaryl or -heteroaryl moiety is reacted with an unsaturated heteroaryl according to Formula (XII) in the presence of catalytic amounts of Pd(OAc)₂ and 1,3-bis diphenylphosphinopropane, in the presence of a suitable base, preferably Cs₂CO₃ or K(AcO), in a reaction-inert polar solvent such as, preferably NMP, DMA, DMF or the like and at an elevated reaction temperature, preferably at 140-150 °C for a certain period of time, preferably about 6-20 hours, more preferably 12-18 hours.



15

Preferably, r is 1. Het may be a unsaturated monocyclic or bicyclic heteroaryl moiety, such as for instance imidazo[1,2-a]pyridinyl, pyrrolyl, thienyl, thiazolyl, imidazolyl, oxazolyl, furanyl, thienyl, benzimidazolyl, benzoxazolyl, benzthiazolyl, benzofuranyl, benzothienyl or indolyl or such as any of the unsaturated radicals in the groups Het¹ and Het² as defined in Formula (I), optionally substituted with one or more radicals selected from the group of Ar¹, Ar¹alkyl, halo, hydroxy, alkyl, piperidinyl, pyrrolyl, thienyl, oxo, alkyloxy, alkyloxyalkyl and alkyloxycarbonyl. Preferably, A is phenyl or pyridinyl.

25

Said reaction has an improved yield over the process according to the prior art (Yutaka Aoyagi et al., *Heterocycles*, 1992, 33, 257 and Sommai Pivsa-Art, *Bull. Chem. Soc. Jpn.*, 1998, 71, 467).

-28-




Also, compounds according to the invention may be converted into an acid addition salt by treatment with an acid, or into a base addition salt by treatment with a base, or conversely, the acid addition salt form may be converted into the free base by treatment with alkali, or the base addition salt may be converted into the free acid by treatment with an acid.

The following examples are intended to illustrate and not to limit the scope of the present invention.


Experimental Part

Hereinafter "RT" means room temperature, "CDI" means 1,1'-carbonyldiimidazole, "DIPE" means diisopropylether, "MIK" means methyl isobutyl keton, "BINAP" means [1,1'-binaphthalene]-2,2'-diylbis[diphenylphosphine], "NMP" means 1-methyl-2-pyrrolidinone, "Pd₂(dba)₃" means tris(dibenzylideneacetone)dipalladium and "DMF" means *N,N*-dimethylformamide.

-29-

Preparation of the intermediate compoundsExample A1a. Preparation of
intermediate compound 1


Et₃N (0.55 mol) was added to a stirring mixture of 7-(phenylmethyl)-1,4-dioxaspiro[4.5]decane (0.5 mol) in toluene (1500ml). 3,5-Bis(trifluoromethyl)benzoyl chloride (0.5 mol) was added over a 1-hour period (exothermic reaction). The mixture was stirred at room temperature for 2 hours, then allowed to stand for the weekend and washed three times with water (500ml, 2x250ml). The organic layer was separated, dried, filtered and the solvent was evaporated. Yielding: 245g (100%). Part of this fraction was crystallized from petroleum ether. The precipitate was filtered off and dried. Yielding: 1.06g of intermediate compound 1.

b. Preparation of
intermediate compound 2

HCl cp (300 ml) was added to a mixture of intermediate compound 1 (0.5 mol) in ethanol (300 ml) and H₂O (300 ml). The reaction mixture was stirred at 60 °C for 20 hours. The precipitate was filtered off, ground, stirred in H₂O, filtered off, washed with petroleum ether and dried. Yielding: 192 g of intermediate compound 2 ((+)-1-[3,5-bis(trifluoromethyl)benzoyl]-2-(phenylmethyl)-4-piperidinone) (89.4%) (mixture of R and S enantiomers).


-30-

c. Preparation of
intermediate compound 3



A mixture of intermediate compound 2 (0.046 mol), 1-(phenylmethyl)piperazine (0.051 mol) and C (0.056 mol) was stirred for 2 hours at 40 °C. The reaction mixture was cooled to room temperature. Ethanol, p.a. (350 ml) was added. BH_3Na (0.138 mol) was added. The resulting reaction mixture was stirred for one hour at room temperature, then for one hour at 50 °C. More BH_3Na (5.2 g) was added and the reaction mixture was stirred for 2 hours at 50 °C. Again, BH_3Na was added and the reaction mixture was stirred overnight at room temperature, then for 2 hours at 50 °C. Water (10 ml) was added. The mixture was stirred for 15 min. CH_2Cl_2 (200 ml) was added and the mixture was stirred for 15 min. The organic phase was separated, dried (MgSO_4), dicalite was added, the mixture was filtered over dicalite, and the filtrate was evaporated. This fraction was separated into (CIS) and (TRANS) by column chromatography over silica gel. The desired (TRANS)-fractions were collected and the solvent was evaporated, giving 14.8 g of residue ((I), 1.06% (CIS)) and 4.9 g of residue ((II), 6% (CIS)). Resolution and purification of those (TRANS)-fractions (± 20 g in total) was obtained by chromatography over stationary phase Chiralcel OD (1900Gr) in Prochrom LC110 35 bar (eluent: hexane/ethanol 90/10). The desired fractions were collected and the solvent was evaporated. Yielding: 9.5 g of intermediate compound 3 (2R-trans)-1-[3,5-bis(trifluoromethyl)benzoyl]-2-(phenylmethyl)-4-[4-(phenylmethyl)-1-piperazinyl]piperidine.

d. Preparation of
intermediate compound 4




A mixture of intermediate compound 3 (0.288 mol) in methanol (700 ml) was hydrogenated at 40 °C with Pd/C, 10% (5 g) as a catalyst. After uptake of H_2 (1 equiv), the catalyst was filtered off and the filtrate was evaporated. Yielding: 141.2 g of

-31-

intermediate compound 4 (+)-(2*R*-trans)-1-[3,5-bis(trifluoromethyl)benzoyl]-2-(phenylmethyl)-4-(1-piperazinyl)piperidine.

Example A2


Preparation of intermediate compound 5



- 5 A mixture of *N*-[(1,1-dimethylethoxy)carbonyl]-L-tyrosine 1,1-dimethylethylcarbonate (0.005 mol), *N,N*-dimethyl-4-pyridinamine (0.006 mol) and Et₃N (0.006 mol) in CH₂Cl₂, p.a. (10ml) was stirred at room temperature. *N*-(ethylcarbonimidoyl)-*N,N*-dimethyl-1,3-propanediamine monohydrochloride (0.006 mol) was added portionwise and was stirred for 45 minutes at room temperature. Then final compound 2 (described in example
- 10 B1.b) (0.005 mol) was added and the reaction mixture was stirred overnight at room temperature. The mixture was washed with H₂O and Na₂CO₃. The separated organic layer was dried, filtered and the solvent was evaporated. The residue was purified over silicagel on a glass filter (eluent : CH₂Cl₂/MeOH 100/0;98/2;96/4;94/6). The purest fractions were collected and the solvent was evaporated. Yield : 1.4g intermediate
- 15 compound 5 (30%).

Example A3

a. Preparation of intermediate compound 6




- A mixture of 7-(hydroxyphenylmethyl)-1,4-dioxo-8-azaspiro[4,5]decane-8-carboxylic acid 1,1-dimethylethyl ester (0.5 mol) and 2-methyl-2-propanol potassium salt (6g) in
- 20 toluene (900ml) was stirred and refluxed for 2h. The mixture was evaporated and the

-32-

residue was stirred up in petrol ether and a little water. The mixture was decanted and the residue was stirred up in DIPE. The precipitate was filtered off and dried. Yielding : 127.4g of intermediate compound 6 (92%).


b. Preparation of
intermediate compound 7



- 5 A mixture of intermediate compound 6 (0.5 mol) in methanol (700ml) was hydrogenated at 50°C overnight with Pd/C, 10% (5g) as a catalyst. After uptake of H₂ (1eq), the catalyst was filtered off and the filtrate was evaporated. The residue was taken up in water and extracted with CH₂Cl₂. The organic layer was dried (MgSO₄), filtered off and evaporated. Yielding : 99g intermediate compound 7 (85%).

10


c. Preparation of
intermediate compound 8



- 15 Et₃N (0.55 mol) was added to a mixture of intermediate compound 7 (0.5 mol) in toluene (1500ml). 3,5-Dimethylbenzoyl chloride (0.5 mol) was added dropwise slowly over a 1-hour period while the temperature was kept below 50°C and while stirring was continued. The mixture was stirred at room temperature overnight, then washed three times with water (500ml, 2x250ml) and separated into its layers. The organic layer was dried (MgSO₄), filtered and the solvent was evaporated. Yielding: 197g (113%). Part of this fraction was dried. Yielding: 0.65g of intermediate compound 8.


-33-

d. Preparation of
intermediate compound 9




A mixture of intermediate compound 8 (0.56 mol) in ethanol (300ml), HCl (300ml) and H₂O (300ml) was stirred at 60°C for 8 hours. The mixture was stirred at room temperature for the weekend. The precipitate was filtered off, taken up in water, filtered off, washed with petroleum ether and dried. Yielding: 140.9g of intermediate compound 9 (88%).

e. Preparation of
intermediate compound 10



A mixture of intermediate compound 9 (0.05 mol) and 1-(phenylmethyl)-piperazine (0.05 mol) in thiophene, 4% solution (2ml) and toluene (500ml) was hydrogenated with Pd/C, 10% (1g) as a catalyst. After uptake of H₂ (1eq), the catalyst was filtered off and the filtrate was evaporated. The residue was purified by column chromatography over silica gel (eluent : CH₂Cl₂/(CH₃OH/NH₃) 99/1). The pure fractions were collected and evaporated. Yielding : 17.07g (71%).. The pure fractions of fraction 1 were collected and evaporated. Yielding : 2.5g of intermediate compound 10 (10%).

f. Preparation of
intermediate compound 11




A mixture of intermediate compound 10 (0.0052 mol) in methanol (100ml) was hydrogenated at 50°C for one night with Pd/C, 10% (1g) as a catalyst. After uptake of H₂ (1eq), the catalyst was filtered off and the filtrate was evaporated. The residue was purified on a glass filter over silica gel (eluent : CH₂Cl₂/(CH₃OH/NH₃) 99/1, 98/2, 97/3,

-34-

96/4 and 95/5). The pure fractions were collected and evaporated. Yielding : 1.7g on intermediate compound 11 (83%).

Example A4

Preparation of intermediate compound 12




- 5 A mixture of final compound 2 (prepared according to B1b) (0.01 mol) and KOH (0.15 mol) in 2-propanol (50 ml) was stirred and refluxed for 18 hours. The solvent was evaporated, then the residue was taken up in H₂O (20 ml) and the mixture was extracted with CH₂Cl₂. The organic layer was washed with NaOH (1N), dried (MgSO₄), filtered and the solvent was evaporated. Yield: 3.25 g of intermediate compound 12 (95%).

10

Preparation of the final compounds

Example B1


a. Preparation of final compound 1



- 15 A mixture of intermediate compound 4 (0.12 mol) and 1-(phenylmethyl)-4-piperidinone (0.12 mol) in methanol (250ml) was hydrogenated at 50°C with Pd/C 10% (3g) as a catalyst in the presence of thiophene solution (2ml). After uptake of H₂ (1 equiv), the catalyst was filtered off and the filtrate was evaporated. The residue was suspended in petroleum ether, filtered off and crystallized from DIPE. Yield : 46g (F1). The filtrate was evaporated. Yield : 37.7g (F2). F1 and F2 were combined and purified by column chromatography over silica gel (eluent : CH₂Cl₂/MeOH 91/9). The product fractions
- 20 were collected and the solvent was evaporated. Yield : 46 g (F3). Part of F3 was crystallized from DIPE. Yield : 0.65 g of final compound 1.

-35-


b. Preparation of final compound 2



A mixture of final compound 1 (0.0074 mol) in methanol (150ml) was hydrogenated with Pd/C 10% (1g) as a catalyst. After uptake of H₂ (1 equiv), the catalyst was filtered off and the filtrate was concentrated. Yield : 4.3g of final compound 2.


5 Example B2

Preparation of final compound 3




- A mixture of final compound 2 (0.0015 mol) and Et₃N (0.1 mol) in CH₂Cl₂ (100ml) was stirred at room temperature. Benzoylchloride (0.0025 mol) was dissolved in CH₂Cl₂ and added dropwise to the reaction mixture. The mixture was stirred for 1 hour at room temperature. NaOH (1N;100ml) was added and the mixture was stirred for 30 minutes at room temperature. The separated aqueous layer was extracted with CH₂Cl₂. The organic layer was washed with H₂O, dried (MgSO₄), filtered and the solvent was evaporated. The residue was purified by column chromatography over silica gel (eluent : CH₂Cl₂/MeOH 100/0;90/10). The desired fractions were collected and the solvent was evaporated. Yield : 0.624g of final compound 3. (61%).

-36-

Example B3a. Preparation of final compound 4

A mixture of 5-methyl-4-isoxazolecarboxylic acid (0.0015 mol) in CH_2Cl_2 (20 ml) and 1,1'-carbonylbis-1*H*-imidazole (0.0015 mol) was stirred for 2 hours at room temperature. Final compound 2 (prepared according to B1.b) (0.001 mol) was added. After stirring overnight, the reaction mixture was washed with diluted NaOH, washed with H_2O , dried, filtered and the solvent evaporated. The residue was purified by column chromatography over silica gel (eluent: CH_2Cl_2 -gradient 0->10% MeOH). The product fractions were collected and the solvent evaporated. The residue was dried. Yield : 0.204 g of final compound 4.

10 b. Preparation of final compound 5



15 A mixture of 3-thiophenecarboxylic acid (0.00188 mol), *N,N*-dimethyl-4-pyridinamine (0.00255 mol) and Et_3N (0.00255 mol) in CH_2Cl_2 (200 ml) was stirred at room temperature. *N,N*-dimethyl-*N'*-(methylcarbonimidoyl)-1,3-propanediamine (0.00255 mol) was added portionwise and the mixture was stirred for one hour at room temperature. A solution of final compound 2 (prepared according to B1b) (0.00188 mol) in CH_2Cl_2 was added dropwise and the reaction mixture was stirred over the weekend at room temperature. The mixture was poured out into 1 g NaOH/water. The layers were separated. The water layer was extracted with CH_2Cl_2 . The separated organic layer was dried (MgSO_4), filtered and the solvent evaporated. The residue was purified by column chromatography over silica gel (eluent: $\text{CH}_2\text{Cl}_2/\text{CH}_3\text{OH}$ from 100/0


20

-37-

to 90/10). The product fractions were collected and the solvent was evaporated. Yield: 0.749 g of final compound 5 (58%).


Example B4

a. Preparation of final compound 6.



- 5 A mixture of final compound 2 (prepared according to B1b) (0.005 mol), 4-(chlorophenylacetyl)-morpholine (0.005 mol) and Na_2CO_3 (0.01 mol) in MIK, p.a. (125 ml) was stirred and refluxed for 18 hours using a water separator. The reaction mixture was washed with water, dried, filtered and the solvent evaporated. The residue was purified over silica gel on a glass filter (eluent: $\text{CH}_2\text{Cl}_2/(\text{CH}_3\text{OH}/\text{NH}_3)$ 95/5). The
- 10 product fractions were collected and the solvent was evaporated. The residue was suspended in DIPE, filtered off and dried. Yield: 1.702 g of final compound 6.

b. Preparation of final compound 7




- A mixture of final compound 2 (prepared according to B1b) (0.0012 mol), 2-(chloromethyl)-1H-benzimidazole (0.0014 mol) and K_2CO_3 (0.0018 mol) in CH_3CN
- 15 (5ml) was stirred and refluxed for 12 hours, then cooled to room temperature and the solvent was evaporated. The residue was taken up in CH_2Cl_2 . The organic layer was washed with H_2O , dried (MgSO_4), filtered and the solvent was evaporated. The residue (0.95g) was purified by column chromatography over silica gel (eluent: $\text{CH}_2\text{Cl}_2/\text{CH}_3\text{OH}/\text{NH}_4\text{OH}$ 90/10/0.5; 15-40 μm). The pure fractions were collected and

-38-


the solvent was evaporated. The residue (0.14g) was crystallized from DIPE. The precipitate was filtered off and dried. Yielding: 0.087g of final compound 7 (10%) (mp. 135°C).

c. Preparation of final compound 8



- 5 A mixture of final compound 2 (prepared according to B1b) (0.005 mol) and 2-(chloromethyl)-6-methyl-3-pyridinol (0.006 mol) was taken up in DMF (50ml). *N*-methyl-*N*-(1-methylethyl)-propanamine (0.02 mol) was added. The reaction mixture was stirred overnight at $\pm 65^{\circ}\text{C}$. The solvent was evaporated. The residue was taken up in CH_2Cl_2 and washed with a diluted NH_3 solution. The separated organic layer was
- 10 dried, filtered and the solvent was evaporated. The residue was purified by column chromatography over silica gel (eluent : $\text{CH}_2\text{Cl}_2/(\text{MeOH}/\text{NH}_3)$ 95/5). The desired fractions were collected and the solvent was evaporated. The residue was suspended in DIPE. The precipitate was filtered off and dried. Yield : 1.423g of final compound 8.

- 15 Example B5
Preparation of final compound 9




A mixture of final compound 2 (prepared according to B1b) (0.003 mol) and 1-methyl-1*H*-pyrrole-2-carboxaldehyde (0.0046 mol) was hydrogenated at 50°C under H_2 with Pd/C 10% (1g) as a catalyst in the presence of thiophene solution (1ml). After uptake of H_2 (1 equiv.), the catalyst was filtered off and the filtrate was evaporated. The residue

-39-

was purified by column chromatography over silica gel (eluent : $\text{CH}_2\text{Cl}_2/(\text{MeOH}/\text{NH}_3)$ 97/3;95/5). The product fractions were collected and the solvent was evaporated. The residue was suspended in petroleumether. Yield : 1.079g of final compound 9.

5 Example B6

Preparation of final compound 10 and 11




$[2\alpha,4\alpha(2R^*,4S^*)]$ = compound 10


$[2\alpha,4\beta(2R^*,4S^*)]$ =compound 11

- A mixture of intermediate compound 2 (prepared according to A1b) (0.005 mol), intermediate compound 11 (prepared according to A3f) (0.005 mol) and $\text{Ti}(\text{O}i\text{Pr})_4$ (3g) in methanol (150ml) was hydrogenated at 50°C under N_2 flow with Pd/C 10% (1g) as a catalyst in the presence of thiophene solution (1ml). After uptake of H_2 (1 equiv), the catalyst was filtered off and the filtrate was evaporated. The residue was taken up in H_2O and CH_2Cl_2 . The mixture was stirred for 10 min and filtered over dicalite. The organic layer was separated, dried, filtered and the solvent was evaporated. The residue was purified by column chromatography over silica gel (eluent: $\text{CH}_2\text{Cl}_2/(\text{CH}_3\text{OH}/\text{NH}_3)$ 97/3). Two fractions were collected and their solvents were evaporated. Yielding: 0.53g compound 10 and 0.4g of final compound 11.

-40-


Example B7Preparation of final
compound 12

- A mixture of final compound 2 (prepared according to B1b) (0.001 mol) in CH_2Cl_2 (50 ml) and CDI (0.0015 mol) was stirred overnight. The reaction mixture was washed with diluted NaOH, washed with H_2O , dried and the solvent was evaporated. The residue was purified by column chromatography over silica gel (Eluent: $\text{CH}_2\text{Cl}_2/\text{CH}_3\text{OH}$ 100/0 and 90/10). The product fractions were collected and the solvent evaporated. Yield : 0.645 g of final compound 12.

Example B8Preparation of final
compound 13

- A mixture of intermediate compound 5 (prepared according to A2) (0.0015 mol) in HCl/2-propanol (5 ml) and methanol (20 ml) was stirred and refluxed for 1 hour. The reaction mixture was crystallized, filtered off and dried. Yield : 0.43 g of final compound 13 (38%)


-41-

Example B9Preparation of final
compound 40

A mixture of final compound 31 (prepared according to B2)(0.065 mmol), 4-pyridinylboronic acid (0.09 mmol), Pd(OAc)₂ (0.015 mmol), 1,3-bis(diphenylphosphino)propane (0.03 mmol), Na₂CO₃, 2M (1 ml) and DME (2 ml) was stirred at 100°C for 16 hours.

- 5 The solvent was evaporated and the residue was taken up in H₂O and extracted with CH₂Cl₂. The organic layer was separated, dried with MgSO₄ and the solvent evaporated. The residue was purified by column chromatography over kromasil (gradient: CH₂Cl₂/CH₃OH 95/5). The desired fractions were collected and the solvent was evaporated. Yield: 1 mg of final compound 40.

10

Example B10Preparation of final
compound 85

A mixture of final compound 83 (prepared according to B2)(0.0004 mol), pyrrolidine (0.0006 mol), Pd₂(dba)₃ (0.00001 mol), BINAP (0.00003 mol) and 2-methyl-2-propanol sodium salt (0.0006 mol) in toluene (5 ml) was stirred at 100°C for 16 hours. The


- 15 The solvent was evaporated and the residue was taken up in H₂O and extracted with CH₂Cl₂. The organic layer was separated, dried with MgSO₄ and the solvent evaporated. The residue was purified by column chromatography over kromasil (gradient:

-42-

CH₂Cl₂/CH₃OH 95/5). The desired fractions were collected and the solvent was evaporated. Yield: 0.119 g of final compound 85.


Example B11

Preparation of final compound 43




- 5 A mixture of final compound 31 (prepared according to B2)(0.065 mmol), imidazo(1,2-a)pyridine (0.09 mmol), Pd(OAc)₂ (0.015 mmol), 1,3-bis(diphenylphosphino)propane (0.03 mmol) and Cs₂CO₃ (0.09 mmol) in NMP (5 ml) was stirred at 140°C for 16 hours. The solvent was evaporated and the residue was taken up in H₂O and extracted with CH₂Cl₂. The organic layer was separated, dried
- 10 with MgSO₄ and the solvent evaporated. The residue was purified by column chromatography over kromasil (gradient: CH₂Cl₂/CH₃OH 95/5). The desired fractions were collected and the solvent was evaporated. The desired fractions were collected and the solvent was evaporated. Yield: 8 mg of final compound 43.

-43-

Example B12Preparation of final
compound 44

- A mixture of compound 31 (prepared according to B2)(0.065 mmol), morpholine (0.2 mmol), Pd(OAc)₂ (0.015 mmol) and 1,3-bis(diphenylphosphino)propane (0.03 mmol) in diglyme (3 ml) under 1 atmosphere CO was stirred at 150°C for 16 hours. The solvent was evaporated and the residue was taken up in H₂O and extracted with CH₂Cl₂. The organic layer was separated, dried with MgSO₄ and the solvent evaporated. The residue was purified by column chromatography over kromasil (gradient: CH₂Cl₂/CH₃OH 95/5). The desired fractions were collected and the solvent was evaporated. The desired fractions were collected and the solvent was evaporated. Yield: 3 mg of final compound 44.

Example B13Preparation of final
compound 144


- A mixture of 4-[(4-acetyloxy)methyl]-1,2,3-thiadiazole-5-carboxylic acid methyl ester (0.001 mol), final compound 2 (prepared according to B1b) (0.002 mol), NaCN (20 mg) in methanol (20ml) was stirred and refluxed for 20 hours. The solvent was evaporated and the residue was purified by column chromatography over silica gel (eluent : CH₂Cl₂/MeOH from 100/0 to 80/20). The desired fractions were collected and the

-44-

solvent was evaporated. The residue was suspended in petroleum ether. The precipitate was filtered off and dried. Yield : 0.110 g of final compound 144.

Example B14


Preparation of final compound 130



- 5 A mixture of final compound 2 (prepared according to B1b) (0.001 mol), glycolaldehyde dimer (0.001 mol) and 3-thiophene boronic acid (0.001 mol) in 2,2,2-trifluoroethanol (5 ml) was stirred at room temperature for 18 hours. This was followed by addition of a solution of K_2CO_3 (10%) and extraction with ethyl acetate. The combined organic layers were dried ($MgSO_4$), filtered and concentrated under vacuum.
- 10 The residue (0.6 g) was purified by chromatography on a silicagel column ($CH_2Cl_2/MeOH/NH_4OH$ 92/08/0.2) and the product fractions were concentrated, providing 0.29 g (47%) of final compound 130.

Example B15

Preparation of final compound 153



- 15 A mixture of intermediate compound 12 (prepared according to A4) (0.00934 mol) and Et_3N (0.02 mol) in CH_2Cl_2 (200 ml) was stirred on an ice bath, then a solution of 4-methyl-1,2,3-thiadiazole-5-carbonyl chloride (0.00943 mol) in CH_2Cl_2 (20 ml) was added dropwise over 15 minutes at $0^\circ C$. The reaction mixture allowed to reach room temperature and was stirred for 1 hour at room temperature, $NaOH$ (20 ml) was added

-45-

and the reaction mixture was stirred for 15 minutes at room temperature. The layers were separated and the aqueous layer was extracted with CH_2Cl_2 . The organic layer was washed with H_2O , dried (MgSO_4), filtered off and the solvent was evaporated. The residue was purified by column chromatography over silicagel (eluent :

- 5 $\text{CH}_2\text{Cl}_2/\text{MeOH}/(\text{MeOH}/\text{NH}_3)$ from 100/0/0 to 90/10/0 to 90/10/0). Two product fractions were collected and each solvent was evaporated. Yield fraction 1: 1.260 g of final compound 153 (22%).

- 10 The compounds exemplified in the following tables were prepared in a manner analogous to one of the foregoing examples B1 to B15.

Analytical data

For a number of compounds, either melting points, LCMS data or optical rotations were recorded.

15

Melting points

If possible, melting points (or ranges) were obtained with a Büchi melting point apparatus B-545. The heating medium is a metal block. The melting of the sample is visually observed by a magnifying lens and a big light contrast. Melting points are

20 measured with a temperature gradient of either 3 or 10 degrees Celsius/minute.

Compound no.	Result (°C)
1	115.9-119.7
2	160.6-163.2
3	149.9-151.7
4	180.5-182.1
5	87.8-121.4
6	87.7-111.2
7	141.0-177.3
8	162.3-164.3
9	122.1-123.8
10	97.0-120.4
11	111.9-125.4
12	66.7-79.0
13	284.5-288.6
14	107.4-116.1
15	188.1-190.3
19	140.3-144.8
22	98.3-119.9
29	142.9-146.5
31	153.1-155.2

-46-

Compound no.	Result (°C)
32	83.3-95.5
33	82.7-98.6
34	80.7-95.5
37	298.1-319.7
38	83.2-110.2
39	279.4-280.9
46	81.3-107.2
49	145.3-149.6
50	92.1-100.7
51	108.9-127.3
52	93.9-104.6
53	156.6-161.0
54	107.6-122.2
55	96.7-106.3
56	171.3-181.5
57	167.4-169.4
58	92.5-102.6
59	79.1-98.2
60	100.5-121.4
62	91.4-120.3
63	86.0-99.4
64	133.6-159.5
65	102.3-105.8
69	108.6-120.6
71	93.5-127.3
72	91.6-103.2
73	100.5-110.5
75	78.8-93.8
76	76.2-93.8
77	273.6-295.2
79	74.3-100.3
80	106.7-126.1
81	85.3-120.6
82	91.9-121.2
83	86.9-102.1
84	92.2-126.1
85	145.4-147.2
88	70.6-108.7
89	96.1-109.4
90	111.9-120.1
91	91.5-108.1
92	100.7-117.9
93	184.1-192.4
98	177.1-180.6
99	65.9-83.0

-47-

Compound no.	Result (°C)
100	76.1-100.1
102	72.9-93.5
103	83.7-100.8
104	105.1-108.5
106	77.2-99.1
108	314.8-335.8
109	95.4-107.7
110	84.6-111.8
111	87.3-109.3
113	252.3-291.7
116	102.8-125.6
117	158.2-160.5
122	177.5°C

LCMS conditions

- The HPLC gradient was supplied by a Waters Alliance HT 2790 system with a column heater set at 40°C. Flow from the column was split to a Waters 996 photodiode array (PDA) detector and a Waters-Micromass ZQ mass spectrometer with an electrospray ionization source operated in positive and negative ionization mode.
- Reversed phase HPLC was carried out on a Xterra MS C18 column (3.5 mm, 4.6 x 100 mm) with a flow rate of 1.6 ml/min. Three mobile phases (mobile phase A 95% 25mM ammoniumacetate + 5% acetonitrile; mobile phase B: acetonitrile; mobile phase C: methanol) were employed to run a gradient condition from 100 % A to 50% B and 50% C in 6.5 min., to 100 % B in 1 min, 100% B for 1 min. and reequilibrate with 100 % A for 1.5 min. An injection volume of 10 mL was used.
- Mass spectra were acquired by scanning from 100 to 1000 in 1 s using a dwell time of 0.1 s. The capillary needle voltage was 3kV and the source temperature was maintained at 140°C. Nitrogen was used as the nebulizer gas. Cone voltage was 10 V for positive ionization mode and 20 V for negative ionization mode. Data acquisition was performed with a Waters-Micromass MassLynx-Openlynx data system.

Compound no.	LCMS MS(MH+)
16	661
18	703
20	711
21	724

-48-

Compound no.	LCMS MS(MH+)
22	701
23	703
24	753
26	809
27	699
28	749
30	654
35	703
36	703
42	756
48	719
61	747
70	693
74	692
94	740
96	703
101	651
105	731
107	691
114	803
115	791
118	859
119	767
124	700
125	673
126	673
127	673
128	737
129	709
130	709
131	693
132	687
133	687
134	687
135	701

-49-

Compound no.	LCMS MS(MH+)
136	677
137	677
138	677
139	709
140	709
141	709
142	709
143	709
144	725
145	681
146	681
147	681
148	651
149	651
150	651
151	677
153	595
154	709
155	709
156	619
157	723
158	745

Optical rotations

Optical rotations were recorded on a polarimeter (Perkin Elmer) at 20°C. Specifics on concentration, wavelength and solvent are given in the table.


5

Compound no.	$[\alpha]$	Wavelength (nm)	Concentration (w/v%)	Solvent
18	-33.77°	365	0.4086	CH ₃ OH
159	-35.56°	365	0.4302	CH ₃ OH
160	-33.66°	365	0.5288	CH ₃ OH
161	-34.75°	365	0.4058	CH ₃ OH
162	-6.72°	436	0.6400	CH ₃ OH
163	-33.2°	365	0.4638	CH ₃ OH
164	-34.1°	365	0.4340	CH ₃ OH









-50-

Compound no.	$[\alpha]$	Wavelength (nm)	Concentration (w/v%)	Solvent
165	-34.43°	365	0.4298	CH ₃ OH
166	-33.95°	365	0.4094	CH ₃ OH
167	-29.91°	365	0.4848	CH ₃ OH
168	-29.12°	365	0.4602	CH ₃ OH
169	-32.32°	365	0.4548	CH ₃ OH
170	-33.3°	365	0.4354	CH ₃ OH
171	-35.06°	365	0.4164	CH ₃ OH
172	-35.84°	365	0.4380	CH ₃ OH
173	-34.53°	365	0.4054	CH ₃ OH












Table 1



















Comp. No.	Exp. No.	Alk ^a	Y	Alk ^b	L	Physical data
2	B1b	cb	cb	cb	H	2R-trans
121	B1b	cb	cb	cb	H	2R-cis
122	B1b	cb	cb	cb	H	2S-trans
123	B1b	cb	cb	cb	H	2S-cis
15	B4b	cb	cb	cb		2R-trans
16	B4a	cb	cb	cb		2R-trans
17	B4c	cb	cb	cb		2R-trans

Comp. No.	Exp. No.	Alk ^a	Y	Alk ^b	L	Physical data
18	B4c	cb	cb	cb		2R-trans
124	B4c	cb	cb	cb		2R-trans
9	B5	-CH ₂ -	cb	cb		2R-trans
20	B4b	-CH ₂ -	cb	cb		2R-trans
8	B4c	-CH ₂ -	cb	cb		2R-trans
7	B4b	-CH ₂ -	cb	cb		2R-trans
21	B4b	-CH ₂ -	cb	cb		B-trans
125	B1a	-CH ₂ -	cb	cb		2R-cis








Comp. No.	Exp. No.	Alk ^a	Y	Alk ^b	L	Physical data
126	B1a	-CH ₂ -	cb	cb		2S-cis
1	B1a	-CH ₂ -	cb	cb		2R-trans
127	B1a	-CH ₂ -	cb	cb		2S-trans
22	B4b	-CH ₂ -	cb	cb		2R-trans
23	B4b	-CH ₂ -	cb	cb		2R-trans
24	B4b	-CH ₂ -	cb	cb		2R-trans







Comp. No.	Exp. No.	Alk ^a	Y	Alk ^b	L	Physical data
25	B4b	-CH ₂ -	cb	cb		B-trans
26	B4b	-CH ₂ -	cb	cb		B-trans
27	B4b	-CH ₂ - CH=CH-	cb	cb		[2B-[2α,4β(E)]]
128	B14		cb	cb		2R-trans
129	B14		cb	cb		2R-trans
130	B14		cb	cb		2R-trans
131	B14		cb	cb		2R-trans

Comp. No.	Exp. No.	Alk ^a	Y	Alk ^b	L	Physical data
28	B4c		cb	cb		B-trans
29	B2	cb	C=O	cb		2R-trans
162	B3b	cb	C=O	cb		2R-trans
30	B2	cb	C=O	cb		2R-trans
3	B2	cb	C=O	cb		2R-trans mp. 142.5°C
132	B2	cb	C=O	cb		2S-trans
133	B2	cb	C=O	cb		2R-cis
134	B2	cb	C=O	cb		2S-cis






Comp. No.	Exp. No.	Alk ^a	Y	Alk ^b	L	Physical data
31	B2	cb	C=O	cb		2R-trans
32	B2	cb	C=O	cb		2R-trans
165	B2	cb	C=O	cb		2R-trans
33	B2	cb	C=O	cb		2R-trans
34	B2	cb	C=O	cb		2R-trans
164	B2	cb	C=O	cb		2R-trans
35	B3b	cb	C=O	cb		2R-trans







Comp. No.	Exp. No.	Alk ^a	Y	Alk ^b	L	Physical data
36	B2	cb	C=O	cb		2R-trans
163	B2	cb	C=O	cb		2R-trans
37	B2	cb	C=O	cb		2R-trans
135	B2	cb	C=O	cb		2R-trans HCl(1:2)
38	B2	cb	C=O	cb		2R-trans
39	B3a	cb	C=O	cb		2R-trans
40	B9	cb	C=O	cb		2R-trans









Comp. No.	Exp. No.	Alk ^a	Y	Alk ^b	L	Physical data
41	B10	cb	C=O	cb		2R-trans
42	B10	cb	C=O	cb		2R-trans
43	B11	cb	C=O	cb		2R-trans
44	B12	cb	C=O	cb		2R-trans
45	B12	cb	C=O	cb		2R-trans
46	B2	cb	C=O	cb		B-trans
47	B2	cb	C=O	cb		2R-trans









Comp. No.	Exp. No.	Alk ^a	Y	Alk ^b	L	Physical data
48	B2	cb	C=O	cb		2R-trans
49	B2	cb	C=O	cb		2R-trans
50	B2	cb	C=O	cb		2R-trans
51	B2	cb	C=O	cb		2R-trans
52	B2	cb	C=O	cb		2R-trans
53	B2	cb	C=O	cb		2R-trans








Comp. No.	Exp. No.	Alk ^a	Y	Alk ^b	L	Physical data
54	B2	cb	C=O	cb		2R-trans
55	B2	cb	C=O	cb		2R-trans
56	B3b	cb	C=O	cb		2R-trans
57	B2	cb	C=O	cb		B-trans
58	B2	cb	C=O	cb		2R-cis

Comp. No.	Exp. No.	Alk ^a	Y	Alk ^p	L	Physical data
59	B2	cb	C=O	cb		B-trans
60	B2	cb	C=O	cb		trans
170	B3b	cb	C=O	cb		2R-trans
61	B2	cb	C=O	cb		2R-trans
62	B2	cb	C=O	cb		2R-trans






Comp. No.	Exp. No.	Alk ^a	Y	Alk ^b	L	Physical data
63	B2	cb	C=O	cb		2R-trans
64	B3a	cb	C=O	cb		2R-trans
65	B2	cb	C=O	cb		2R-trans
66	B2	cb	C=O	cb		B-trans
67	B3b	cb	C=O	cb		2R-trans
68	B2	cb	C=O	cb		2R-trans







Comp. No.	Exp. No.	Alk ^a	Y	Alk ^b	L	Physical data
69	B3a	cb	C=O	cb		2R-trans
5	B3b	cb	C=O	cb		2R-trans
70	B2	cb	C=O	cb		2R-trans
161	B2	cb	C=O	cb		2R-trans
71	B3a	cb	C=O	cb		2R-trans
136	B3b	cb	C=O	cb		2S-trans
137	B3b	cb	C=O	cb		2R-cis
138	B3b	cb	C=O	cb		2S-cis










Comp. No.	Exp. No.	Alk ^a	Y	Alk ^b	L	Physical data
72	B3a	cb	C=O	cb		2R-trans
12	B7	cb	C=O	cb		2R-trans
73	B2	cb	C=O	cb		2R-trans
19	B2	cb	C=O	cb		2R-trans
74	B2	cb	C=O	cb		2R-trans
75	B2	cb	C=O	cb		2R-trans
4	B3a	cb	C=O	cb		2R-trans
76	B3a	cb	C=O	cb		2R-trans












Comp. No.	Exp. No.	Alk ^a	Y	Alk ^b	L	Physical data
77	B2	cb	C=O	cb		2R-trans m.p. 119.6 °C
139	B2	cb	C=O	cb		2R-cis
140	B2	cb	C=O	cb		2S-cis
141	B2	cb	C=O	cb		2S-trans
78	B2	cb	C=O	cb		2R-trans; HCl(1:2); H ₂ O(1:1)
142	B2	cb	C=O	cb		2R-trans; succinate (1:2)
143	B2	cb	C=O	cb		2R-trans; malonate (1:2)











Comp. No.	Exp. No.	Alk ^a	Y	Alk ^b	L	Physical data
144	B13	cb	C=O	cb		2R-trans
120	B3b	cb	C=O	cb		2R-trans
79	B2	cb	C=O	cb		2R-trans
166	B3b	cb	C=O	cb		2R-trans
80	B2	cb	C=O	cb		2R-trans
81	B3b	cb	C=O	cb		2R-trans
82	B3b	cb	C=O	cb		2R-trans















Comp. No.	Exp. No.	Alk ^a	Y	Alk ^b	L	Physical data
83	B2	cb	C=O	cb		2R-trans
14	B2	cb	C=O	cb		2R-trans
84	B3b	cb	C=O	cb		2R-trans
85	B10	cb	C=O	cb		2R-trans
86	B9	cb	C=O	cb		2R-trans













Comp. No.	Exp. No.	Alk ^a	Y	Alk ^b	L	Physical data
87	B9	cb	C=O	cb		2R-trans
88	B8	cb	C=O	cb		2R-trans
89	B3b	cb	C=O	cb		2R-trans
90	B3b	cb	C=O	cb		[2R- [2 α ,4 β (S)]]
91	B8	cb	C=O	cb		[2R- [2 α ,4 β (S)]]
92	B2	cb	C=O	cb		2R-trans





Comp. No.	Exp. No.	Alk ^a	Y	Alk ^b	L	Physical data
93	B3b	cb	C=O	cb		2R-trans
94	B3b	cb	C=O	cb		B-trans
169	B3b	cb	C=O	cb		2R-trans
96	B2	cb	C=O	cb		B-trans
145	B3b	cb	C=O	cb		2S-trans
146	B3b	cb	C=O	cb		2R-cis
147	B3b	cb	C=O	cb		2S-cis
173		cb	C=O	cb		
97	B4c	-CH ₂ -	C=O	cb		2R-trans

Comp. No.	Exp. No.	Alk ^a	Y	Alk ^b	L	Physical data
98	B2	cb	C=O	-CH ₂ -	-H	2R-trans
99	B2	cb	C=O		-H	2R-trans
159	B2	cb	C=O		-H	2R-trans
167	B3b	cb	C=O			2R-trans
160	B2	cb	C=O		-H	2R-trans
100	B2	cb	C=O		-H	2R-trans
101	B2	cb	C=O		-H	2R-trans
148	B2	cb	C=O		-H	2S-trans
149	B2	cb	C=O		-H	2R-cis
150	B2	cb	C=O		-H	2S-cis
171	B3b	cb	C=O		-H	2R-trans

Comp. No.	Exp. No.	Alk ^a	Y	Alk ^b	L	Physical data
172	B3b	cb	C=O		-H	2R-trans
102	B2	cb	C=O		-H	2R-trans
151	B2	cb	C=O		-H	2R-trans
103	B2	cb	C=O		-H	2R-trans
104	B2	cb	C=O	-CH ₂ -		2R-trans
105	B2	cb	C=O	-CH ₂ -		2R-trans
106	B2	cb	C=O	-CH ₂ -		2R-trans
107	B3b	cb	C=O	-CH ₂ -		2R-trans
13	B8	cb	C=O			2R-trans, HCl(1:3); H ₂ O(1:1)

Comp. No.	Exp. No.	Alk ^a	Y	Alk ^b	L	Physical data
108	B2	cb	C=O			2R-trans HCl(1:2) H ₂ O(1:1)
109	B2	cb	C=O			2R-trans
110	B3b	cb	C=O			[2R-[2α,4β(E)]]
111	B2	cb	C=O			2R-trans
112	B2	cb	C=O			2R-trans
152	B4c	cb	C=O			B-trans
113	B4c	cb	C=O			B-trans HCl(1:3) H ₂ O(1:3)


Comp. No.	Exp. No.	Alk ^a	Y	Alk ^b	L	Physical data
114	B4b	cb	C=O			B-trans
115	B3b	cb	C=O			B-trans
116	B4c	cb	C=O			2R-trans
6	B4a		C=O	cb		2R-trans
117	B2		C=O	cb		2R-trans
168	B2	cb		cb		2R-trans

Comp. No.	Exp. No.	Alk ^a	Y	Alk ^b	L	Physical data
118	B2	cb		cb		B-trans
119	B2	cb		cb		B-trans

cb = Covalent Bond

-75-

Table 2:




Co No.	Exp. No.	R¹	Alkᵃ	Y	Alkᵇ	L	Physical data
10	B6		cb	C=O	cb		[2α,4α(2R*,4S*)]
11	B6		cb	C=O	cb		[2α,4β(2R*,4S*)]

cb = Covalent Bond

5

Table 3:




Co No.	Exp. No.	Alkᵃ	Y	Alkᵇ	L	Physical data
153	B15	cb	C=O	cb		2R-trans

cb = Covalent Bond

-76-

Table 4:




Co No.	Exp No.	Alk ^a	Y	Alk ^b	L	Physical data
154	B1a	-CH ₂ -	cb	cb		2R-cis
155	B1a	-CH ₂ -	cb	cb		2R-trans
156	B1b	cb	cb	cb	-H	2R-trans
157	B2	cb	C=O	cb		2R-trans
158	B2	cb	C=O	cb		2R-trans

cb = Covalent Bond


5

Table 5:



Co No.	Exp No.	Alk ^a	Y	Alk ^b	L	Physical data
175	B1b	cb	cb	cb	H	cis
174	B1a	-CH ₂ -	cb	cb		
176	B2	cb	C=O	cb		cis

-77-

Co No.	Exp No.	Alk ^a	Y	Alk ^b	L	Physical data
177	B2	cb	C=O	cb		cis

-78-

C. Pharmacological exampleExample C.1 : Binding experiment for h-NK₁, h-NK₂ and h-NK₃ receptors

The compounds according to the invention were investigated for interaction with various neurotransmitter receptors, ion channels and transporter binding sites using the radioligand binding technique. Membranes from tissue homogenates or from cells, expressing the receptor or transporter of interests, were incubated with a radioactively labelled substance ($[^3\text{H}]$ - or $[^{125}\text{I}]$ ligand) to label a particular receptor. Specific receptor binding of the radioligand was distinguished from the non-specific membrane labelling by selectively inhibiting the receptor labelling with an unlabelled drug (the blank), known to compete with the radioligand for binding to the receptor sites. Following incubation, labelled membranes were harvested and rinsed with excessive cold buffer to remove non-bound radioactivity by rapid filtration under suction. Membrane bound radioactivity was counted in a scintillation counter and results were expressed in counts per minute (cpm).

The compounds were dissolved in DMSO and tested at 10 concentrations ranging from 10^{-10} to 10^{-5} M.

The ability of the compounds according to the invention to displace $[^3\text{H}]$ -Substance P from cloned human h-NK₁ receptors expressed in CHO cells, to displace $[^3\text{H}]$ -SR-48968 from cloned human h-NK₂ receptors expressed in Sf9 cells, and to displace $[^3\text{H}]$ -SR-142801 from cloned human h-NK₃ receptors expressed in CHO cells was evaluated.

The pIC₅₀ data for the h-NK₁, h-NK₂ and h-NK₃ receptor testing for a representative selection of compounds are presented in Table 3.

All selected compounds show (sub)nanomolar affinity for the h-NK₁ receptor most of them with more than 100-fold selectivity towards the h-NK₂ and h-NK₃ receptors.

Example C.2 : Signal transduction

This test evaluates in vitro functional NK₁ antagonistic activity. For the measurements of intracellular Ca^{++} concentrations the cells were grown on 96-well (black wall/transparent bottom) plates from Costar for 2 days until they reached confluence. The cells were loaded with 2 μM Fluo3 in DMEM containing 0.1% BSA and 2.5 mM probenecid for 1 h at 37°C. They were washed 3x with a Krebs buffer (140 mM NaCl, 1 mM $\text{MgCl}_2 \cdot 6\text{H}_2\text{O}$, 5 mM KCl, 10 mM glucose, 5 mM HEPES; 1.25 mM CaCl_2 ; pH 7.4) containing 2.5 mM probenecid and 0.1 % BSA (Ca^{++} -buffer). The cells were

preincubated with a concentration range of antagonists for 20 min at RT and Ca^{++} -signals after addition of the agonists were measured in a Fluorescence Image Plate Reader (FLIPR from Molecular Devices, Crawley, England). The peak of the Ca^{++} -transient was considered as the relevant signal and the mean values of corresponding wells were analysed as described below.

The sigmoidal dose response curves were analysed by computerised curve-fitting, using the GraphPad Program. The EC_{50} -value of a compound is the effective dose showing 50 % of maximal effect. For mean curves the response to the agonist with the highest potency was normalised to 100 %. For antagonist responses the IC_{50} -value was calculated using non-linear regression.

Table

Co No.	h-NK ₁ pIC ₅₀	h-NK ₂ pIC ₅₀	h-NK ₃ pIC ₅₀
2	8.6	5.8	5.2
3	8.9	6.3	6.6
4	8.8	5.2	6.7
5	10.0	6.1	6.3
6	9.0	-	-
7	8.1	6.0	6.0
8	9.2	-	-
9	8.9	6.2	6.3
10	7.3	6.4	6.2
11	7.4	6.2	6.6
12	9.1	6.0	6.1
13	8.9	6.2	6.0
16	9.0	6.3	6.8
22	9.4	6.2	6.5
26	7.4	6.0	6.0
32	8.8	6.2	6.8
36	9.0	6.1	6.1
37	8.4	6.3	6.6
39	9.1	6.0	6.0
42	8.6	-	-
45	9.5	-	-
51	8.9	6.2	6.4

-80-

Co No.	h-NK ₁ pIC ₅₀	h-NK ₂ pIC ₅₀	h-NK ₃ pIC ₅₀
56	9.0	6.3	6.7
62	9.2	6.4	6.6
64	8.1	6.4	6.4
65	8.4	6.2	6.6
77	9.0	6.1	5.6
78	9.1	6.4	6.0
79	8.2	6.5	6.4
80	9.3	6.1	6.6
85	8.5	-	-
89	8.6	6.2	6.2
90	7.5	6.5	6.9
97	9.5	6.3	6.4
102	9.0	-	-
104	9.2	5.8	5.8
106	9.0	6.0	6.3
108	8.8	-	-
110	10.0	-	-
113	9.0	6.4	6.4
116	8.6	6.1	6.8
119	7.6	6.0	6.0
132	8.0	5.7	5.5
133	8.4	5.9	6.1
134	7.7	5.6	<5
139	8.8	6.1	6.5
140	8.5	5.4	5.3
141	8.1	5.4	5.4
142	8.9	6.2	6.6
143	9.0	6.1	6.3
144	?	5.9	6.2
151	9.4	6.2	6.4

D. Composition examples

"Active ingredient" (A.I.) as used throughout these examples relates to a compound of Formula (I), the pharmaceutically acceptable acid or base addition salts thereof, the stereochemically isomeric forms thereof, the *N*-oxide form thereof and prodrugs thereof.

5

Example D.1 : ORAL DROPS

500 Grams of the A.I. was dissolved in 0.5 l of 2-hydroxypropanoic acid and 1.5 l of the polyethylene glycol at 60~80°C. After cooling to 30~40°C there were added 35 l of polyethylene glycol and the mixture was stirred well. Then there was added a solution of 10 1750 grams of sodium saccharin in 2.5 l of purified water and while stirring there were added 2.5 l of cocoa flavor and polyethylene glycol q.s. to a volume of 50 l, providing an oral drop solution comprising 10 mg/ml of A.I. The resulting solution was filled into suitable containers.

15 Example D.2 : ORAL SOLUTION

9 Grams of methyl 4-hydroxybenzoate and 1 gram of propyl 4-hydroxybenzoate were dissolved in 4 l of boiling purified water. In 3 l of this solution were dissolved first 10 grams of 2,3-dihydroxybutanedioic acid and thereafter 20 grams of the A.I. The latter solution was combined with the remaining part of the former solution and 12 l 1,2,3- 20 propanetriol and 3 l of sorbitol 70% solution were added thereto. 40 Grams of sodium saccharin were dissolved in 0.5 l of water and 2 ml of raspberry and 2 ml of gooseberry essence were added. The latter solution was combined with the former, water was added q.s. to a volume of 20 l providing an oral solution comprising 5 mg of the active ingredient per teaspoonful (5 ml). The resulting solution was filled in suitable containers.

25

Example D.3 : FILM-COATED TABLETSPreparation of tablet core

A mixture of 100 grams of the A.I., 570 grams lactose and 200 grams starch was mixed well and thereafter humidified with a solution of 5 grams sodium dodecyl sulfate and 10 30 grams polyvinylpyrrolidone in about 200 ml of water. The wet powder mixture was sieved, dried and sieved again. Then there was added 100 grams microcrystalline cellulose and 15 grams hydrogenated vegetable oil. The whole was mixed well and compressed into tablets, giving 10.000 tablets, each containing 10 mg of the active ingredient.

35

Coating

-82-


To a solution of 10 grams methyl cellulose in 75 ml of denaturated ethanol there was added a solution of 5 grams of ethyl cellulose in 150 ml of dichloromethane. Then there were added 75 ml of dichloromethane and 2.5 ml 1,2,3-propanetriol. 10 Grams of polyethylene glycol was molten and dissolved in 75 ml of dichloromethane. The latter
5 solution was added to the former and then there were added 2.5 grams of magnesium octadecanoate, 5 grams of polyvinylpyrrolidone and 30 ml of concentrated color suspension and the whole was homogenated. The tablet cores were coated with the thus obtained mixture in a coating apparatus.

10 Example D.4 : INJECTABLE SOLUTION

1.8 Grams methyl 4-hydroxybenzoate and 0.2 grams propyl 4-hydroxybenzoate were dissolved in about 0.5 l of boiling water for injection. After cooling to about 50°C there were added while stirring 4 grams lactic acid, 0.05 grams propylene glycol and 4 grams of the A.I.. The solution was cooled to room temperature and supplemented with water
15 for injection q.s. ad 1 l, giving a solution comprising 4 mg/ml of A.I.. The solution was sterilized by filtration and filled in sterile containers.

CLAIMS

1. A compound according to the general Formula (I)



5

the pharmaceutically acceptable acid or base addition salts thereof, the stereochemically isomeric forms thereof, the *N*-oxide form thereof and prodrugs thereof, wherein :

- 10 n is an integer, equal to 0, 1 or 2 ;
 m is an integer, equal to 1 or 2, provided that if m is 2, then n is 1 ;
 p is an integer equal to 1 or 2 ;
 Q is O or NR^3 ;
 X is a covalent bond or a bivalent radical of formula -O-, -S- or $-NR^3$ - ;
each R^3 independently from each other, is hydrogen or alkyl ;
15 each R^1 independently from each other, is selected from the group of Ar^1 , Ar^1 -alkyl and $di(Ar^1)$ -alkyl ;
 q is an integer equal to 0 or 1 ;
 R^2 is alkyl, Ar^2 , Ar^2 -alkyl, Het^1 or Het^1 -alkyl ;
 Y is a covalent bond or a bivalent radical of formula $-C(=O)-$ or $-SO_2-$;
20 each Alk represents, independently from each other, a covalent bond; a bivalent straight or branched, saturated or unsaturated hydrocarbon radical having from 1 to 6 carbon atoms ; or a cyclic saturated or unsaturated hydrocarbon radical having from 3 to 6 carbon atoms ; each radical optionally substituted on one or more carbon atoms with one or more
25 alkyl, phenyl, halo, cyano, hydroxy, formyl and amino radicals ;
 L is selected from the group of hydrogen, alkyloxy, Ar^3 -oxy, alkyloxycarbonyl, mono- and $di(alkyl)$ amino, mono- and $di(Ar^3)$ amino, mono- and $di(alkyloxycarbonyl)$ amino, Ar^3 , Ar^3 -carbonyl, Het^2 and Het^2 -carbonyl ;
30 Ar^1 is phenyl, optionally substituted with 1, 2 or 3 substituents each independently from each other selected from the group of halo, alkyl, cyano, aminocarbonyl and alkyloxy ;
 Ar^2 is naphthalenyl or phenyl, each optionally substituted with 1, 2 or 3

- substituents, each independently from each other, selected from the group of halo, nitro, amino, mono- and di(alkyl)amino, cyano, alkyl, hydroxy, alkyloxy, carboxyl, alkyloxycarbonyl, aminocarbonyl and mono- and di(alkyl)aminocarbonyl ;
- 5 Ar^3 is naphthalenyl or phenyl, optionally substituted with 1, 2 or 3 substituents each independently from each other selected from the group of alkyloxy, alkyl, halo, hydroxy, pyridinyl, morpholinyl, pyrrolidinyl, imidazo[1,2-*a*]pyridinyl, morpholinylcarbonyl, pyrrolidinylcarbonyl, amino and cyano;
- 10 Het^1 is a monocyclic heterocyclic radical selected from the group of pyrrolyl, pyrazolyl, imidazolyl, furanyl, thienyl, oxazolyl, isoxazolyl, thiazolyl, thiadiazolyl, isothiazolyl, pyridinyl, pyrimidinyl, pyrazinyl and pyridazinyl ; or a bicyclic heterocyclic radical selected from the group of quinolinyl, quinoxalinyl, indolyl, benzimidazolyl, benzoxazolyl, benzisoxazolyl, benzothiazolyl, benzisothiazolyl, benzofuranyl and benzothienyl ; each heterocyclic radical may optionally be substituted on
- 15 any atom by a radical selected from the group of halo and alkyl ;
- Het^2 is a monocyclic heterocyclic radical selected from the group of pyrrolidinyl, dioxolyl, imidazolidinyl, pyrazolidinyl, piperidinyl, morpholinyl, dithianyl, thiomorpholinyl, piperazinyl, imidazolidinyl, tetrahydrofuranyl, 2H-pyrrolyl, pyrrolinyl, imidazolinyl, pyrazolinyl, pyrrolyl, imidazolyl, pyrazolyl, triazolyl, furanyl, thienyl, oxazolyl, isoxazolyl, thiazolyl, thiadiazolyl, isothiazolyl, pyridinyl, pyrimidinyl, pyrazinyl, pyridazinyl and triazinyl ; or a bicyclic heterocyclic radical
- 20 selected from the group of benzopiperidinyl, quinolinyl, quinoxalinyl, indolyl, isoindolyl, chromenyl, benzimidazolyl, imidazo[1,2-*a*]pyridinyl, benzoxazolyl, benzisoxazolyl, benzothiazolyl, benzisothiazolyl, benzofuranyl and benzothienyl ; each radical optionally substituted with one or more radicals selected from the group of Ar^1 , Ar^1 alkyl, halo,
- 25 hydroxy, alkyl, piperidinyl, pyrrolyl, thienyl, oxo, alkyloxy, alkyloxyalkyl and alkyloxycarbonyl ; and
- 30 alkyl is a straight or branched saturated hydrocarbon radical having from 1 to 6 carbon atoms or a cyclic saturated hydrocarbon radicals having from 3 to 6 carbon atoms ; optionally substituted on one or more carbon atoms
- 35 with one or more radicals selected from the group of phenyl, halo, cyano, oxo, hydroxy, formyl and amino radicals.

2. A compound according to claim 1, characterized in that


-85-

	n	is 1 ;
	m	is 1 ;
	p	is 1 ;
	Q	is O ;
5	X	is a covalent bond ;
	each R ¹	is Ar ¹ or Ar ¹ -alkyl ;
	q	is 0 or 1 ;
	R ²	is Ar ² ;
	Y	is a covalent bond or a bivalent radical of formula -C(=O)- or -SO ₂ - ;
10	each Alk	represents, independently from each other, a covalent bond; a bivalent straight or branched, saturated or unsaturated hydrocarbon radical having from 1 to 6 carbon atoms ; or a cyclic saturated or unsaturated hydrocarbon radical having from 3 to 6 carbon atoms ; each radical optionally substituted on one or more carbon atoms with one or more
15		phenyl, halo, cyano, hydroxy, formyl and amino radicals ;
	L	is selected from the group of hydrogen, alkyloxy, Ar ³ -oxy, alkyloxy-carbonyl, mono- and di(alkyl)amino, mono-and di(Ar ³)amino, Ar ³ and Het ² ;
	Ar ¹	is phenyl, optionally substituted with 1, 2 or 3 alkyl radicals ;
20	Ar ²	is phenyl, optionally substituted with 1, 2 or 3 alkyl radicals ;
	Ar ³	is phenyl, optionally substituted with 1, 2 or 3 substituents each independently from each other selected from the group of alkyloxy, alkyl, halo, hydroxy, pyridinyl, morpholinyl, pyrrolidinyl, imidazo[1,2- <i>a</i>]pyridinyl, morpholinylcarbonyl, pyrrolidinylcarbonyl, amino and
25		cyano ;
	Het ²	is a monocyclic heterocyclic radical selected from the group of pyrrolidinyl, piperidinyl, morpholinyl, pyrrolyl, imidazolyl, pyrazolyl, furanyl, thienyl, isoxazolyl, thiazolyl, thiadiazolyl, pyridinyl, pyrimidinyl, pyrazinyl, and pyridazinyl ; or a bicyclic heterocyclic
30		radical selected from the group of benzopiperidinyl, quinolinyl, quinoxalinyl, indolyl, chromenyl and benzimidazolyl ; each radical optionally substituted with one or more radicals selected from the group of Ar ¹ , Ar ¹ alkyl, halo, hydroxy, alkyl, piperidinyl, pyrrolyl, thienyl, oxo and alkyloxycarbonyl ; and
35	alkyl	is a straight hydrocarbon radical having 1 to 6 carbon atoms, optionally substituted with one or more halo radicals .


-86-

3. A compound according to any of claims 1-2, characterized in that R¹ is Ar¹methyl and attached to the 2-position or R¹ is Ar¹ and attached to the 3-position.
4. A compound according to any of claims 1-3, characterized in that the R²-X-C(=Q)-
5 moiety is 3,5-di-(trifluoromethyl) phenylcarbonyl.
5. A compound according to any one of claims 1-4 for use as a medicine.
6. The use of a compound according to any one of claims 1-4 for the manufacture of a
10 medicament for treating neurokinin mediated conditions.
7. The use of a compound according to claim 6 for the manufacture of a medicament for treating emesis, depression, anxiety disorders, pain, pancreatitis and irritable bowel syndrome (IBS).
15
8. A pharmaceutical composition comprising a pharmaceutically acceptable carrier and, as active ingredient, a therapeutically effective amount of a compound according to any one of claims 1- 4.
9. A pharmaceutical composition according to claim 8, characterized in that it is in a
20 form suitable to be orally administered.
10. A process for the preparation of a composition as claimed in any one of claims 1-4, characterized in that a pharmaceutically acceptable carrier is intimately mixed with a
25 therapeutically effective amount of a compound as claimed in any one of claims 1-4.
11. A process for the preparation of a compound according to Formula (I), more specifically according to Formula (I^a), Formula (I^b) or Formula (I^c), characterized in
30 that
 - a) a final compound according to Formula (I) is obtained by reductive *N*-alkylation of an intermediate according to Formula (II) wherein R¹, R², X, Q, m, n and p are defined as in Formula (I), with an *N*-substituted piperidinon of Formula (III) wherein R¹, Alk, Y, L and q are defined as in Formula (I), in a reaction-inert solvent
35 and in the presence of a reducing agent ; or


-87-



b) a final compound according to Formula (I^a) is obtained by acylation of a final compound of Formula (I') wherein R¹, R², X, Q, m, n, p and q are defined as in Formula (I), with an acyl compound of Formula (V) wherein Alk and L are defined as in Formula (I) and W¹ is a leaving group, in a reaction-inert solvent and in the presence of a base ; or




c) a final compound according to Formula (I^a) is obtained by a base-catalyzed nucleophilic addition reaction of a final compound of Formula (I') wherein R¹, R², X, Q, m, n, p and q are defined as in Formula (I), with a carboxylic acid of Formula (VI) wherein Alk and L are defined as in Formula (I), or its ester, in a reaction-inert solvent and in the presence of a base ; or




d) a final compound according to Formula (I^b) is obtained by a base-catalyzed nucleophilic addition reaction of a final compound of Formula (I') wherein R¹, R², X, Q, m, n, p and q are defined as in Formula (I), with a compound of Formula (VIII) wherein Alk and L are defined as in Formula (I) and W² is a leaving group, in a reaction-inert solvent and in the presence of a base ; or

-88-





e) a final compound according to Formula (I^c) is obtained by reductive amination/alkylation of a final compound of Formula (I') wherein R¹, R², X, Q, m, n, p and q are defined as in Formula (I) with a compound of Formula (VIII) wherein Alk and L are defined as in Formula (I) and W³ is a leaving group, in a reaction-inert solvent and in the presence of a base ; or



f) a final compound according to Formula (I) is obtained by converting compounds according to Formula (I) into each other following art-known transformation reactions ; and further, converting compounds according to Formula (I) into an acid addition salt by treatment with an acid, or into a base addition salt by treatment with a base, or conversely, the acid addition salt form may be converted into the free base by treatment with alkali, or the base addition salt may be converted into the free acid by treatment with an acid ; and by preparing the *N*-oxide and/or stereochemically isomeric forms thereof.

12. A process for the preparation of a compound according to Formula (XIII), characterized in that a compound according to Formula (XI), wherein A is an aryl or heteroaryl, Z may be any moiety, preferably a moiety Z¹ as defined below wherein each variable is defined as in Formula (I), Hal is an halogen and r is an integer ranging from 1 to a number equal to the number of available carbon atoms in the aryl or heteroaryl-moiety A, is reacted with an unsaturated heteroaryl Het according to Formula (XII) in the presence of catalytic amounts of Pd(OAc)₂ and 1,3-bis diphenylphosphinopropane, in the presence of a suitable base, preferably Cs₂CO₃ or K(AcO) and in a reaction-inert polar solvent.

-89-



13. A process according to claim 12, wherein Hal is bromo or iodo, A is phenyl or pyridinyl, Z is Z¹ and Het is selected from the group of imidazo[1,2-a]pyridinyl, pyrrolyl and thienyl.

INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 03/50697

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C07D211/58 C07D413/14 C07D409/14 C07D401/14 C07D471/04
 C07D405/14 C07D417/14

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07D

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the International search (name of data base and, where practical, search terms used)

EPO-Internal, CHEM ABS Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 97/16440 A (JANSSEN PHARMACEUTICA NV ; JANSSENS FRANS EDUARD (BE); SOMMEN FRANCOIS) 9 May 1997 (1997-05-09) cited in the application claims	1,6,8

☐ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the International filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the International filing date but later than the priority date claimed

"T" later document published after the International filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&" document member of the same patent family

Date of the actual completion of the International search

28 January 2004

Date of mailing of the International search report

10/02/2004

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
 NL - 2280 HV Rijswijk
 Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
 Fax: (+31-70) 340-3016

Authorized officer

Diederer, J

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/EP 03/50697

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9716440	A	09-05-1997	
		AT 188691 T	15-01-2000
		AU 704155 B2	15-04-1999
		AU 7493296 A	22-05-1997
		BR 9611184 A	30-03-1999
		CA 2234096 A1	09-05-1997
		CN 1438220 A	27-08-2003
		CN 1205699 A , B	20-01-1999
		CY 2177 A	23-08-2002
		CZ 9801322 A3	16-09-1998
		DE 69606196 D1	17-02-2000
		DE 69606196 T2	21-09-2000
		EA 980404 A1	29-10-1998
		WO 9716440 A1	09-05-1997
		EP 0862566 A1	09-09-1998
		ES 2143238 T3	01-05-2000
		GR 3033154 T3	31-08-2000
		HR 960507 A1	28-02-1998
		HU 9802985 A2	28-10-1999
		IL 123962 A	11-01-2001
		JP 11514634 T	14-12-1999
		JP 3073238 B2	07-08-2000
		NO 981534 A	24-06-1998
		NZ 321575 A	28-05-1999
		PL 327406 A1	07-12-1998
		PT 862566 T	30-06-2000
		SI 862566 T1	30-04-2000
		TR 9800756 T2	21-07-1998
		TW 460473 B	21-10-2001
		US 6197772 B1	06-03-2001
		US 6521621 B1	18-02-2003
		US RE37886 E1	15-10-2002
		ZA 9609090 A	29-04-1998

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☒ **BLACK BORDERS**
- ☐ **IMAGE CUT OFF AT TOP, BOTTOM OR SIDES**
- ☐ **FADED TEXT OR DRAWING**
- ☐ **BLURRED OR ILLEGIBLE TEXT OR DRAWING**
- ☐ **SKEWED/SLANTED IMAGES**
- ☐ **COLOR OR BLACK AND WHITE PHOTOGRAPHS**
- ☐ **GRAY SCALE DOCUMENTS**
- ☐ **LINES OR MARKS ON ORIGINAL DOCUMENT**
- ☐ **REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY**
- ☐ **OTHER:** _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.